arXiv:1207.7336v1 [math.AP] 31 Jul 2012

ENERGY DECAY RATES FOR SOLUTIONS OF THE WAVE EQUATION WITH SUPERLINEAR DAMPING IN EXTERIOR DOMAIN

M. DAOULATLI

Abstract. In this paper we study the behavior of the energy of solutions of the wave equation with localized superlinear damping in exterior domain.

1. Introduction and Statement of the results

Let O be a compact domain of $\mathbb{R}^d$ $(d \geq 1)$ with C^∞ boundary Γ and $\Omega = \mathbb{R}^d \backslash O$. Consider the following wave equatin with localized linear damping

$$\begin{cases} \partial_t^2 u - \Delta u + a(x) \left|\partial_t u\right|^{r-1} \partial_t u = 0 & \text{in } \mathbb{R}_+ \times \Omega, \\ u = 0 & \text{on } \mathbb{R}_+ \times \Gamma, \\ u(0,x) = u_0 \quad \text{and} \quad \partial_t u(0,x) = u_1. & \end{cases} \tag{1.1}$$

Here Δ denotes the Laplace operator in the space variables. $a(x)$ is a nonnegative function in $L^\infty(\Omega)$. Throughout this paper we assume that $1 < r \leq 1 + \frac{2}{d}$. We note that when $d = 1$ and O is not empty , then $\Omega = (-\infty, \alpha)$ or $(\alpha, +\infty)$ for some $\alpha \in \mathbb{R}$.

The existence and uniqueness of global solutions to the problem (1.1) is standard (see [23]). If (u_0, u_1) is in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$, then the system (1.1), admits a unique solution u in the class

$$u \in C^0\left(\mathbb{R}_+, H_0^1(\Omega) \cap H^2(\Omega)\right) \cap C^1\left(\mathbb{R}_+, H_0^1(\Omega)\right).$$

Let us consider the energy at instant t defined by

$$E_u(t) = \frac{1}{2} \int_\Omega \left(\left|\nabla u(t,x)\right|^2 + \left|\partial_t u(t,x)\right|^2 \right) dx.$$

The energy functional satisfies the following identity

$$E_u(T) + \int_0^T \int_\Omega a(x) \left|\partial_t u\right|^{r+1} dx dt = E_u(0), \tag{1.2}$$

for every $T \geq 0$. Moreover we have

$$\begin{aligned} &\left\|\nabla \partial_t u\right\|^2_{L^\infty(\mathbb{R}_+, L^2(\Omega))} + \left\|\partial_t^2 u\right\|^2_{L^\infty(\mathbb{R}_+, L^2(\Omega))} \\ &\leq 2\left(1 + \|a\|_{L^\infty}\right) \left(\|u_0\|^2_{H^2} + \|u_1\|^2_{H^1} + \|u_1\|^{2r}_{H^1} \right) \end{aligned} \tag{1.3}$$

First we quote results when the damping is linear.

Zuazua [36], Nakao [27], Dehman-Lebeau-Zuazua [13], Aloui-Ibrahim-Nakanishi [1] and Joly and Camille [19] have considered the problem for the Klein-Gordon type wave equations

---

Date: August 1, 2012.

2000 *Mathematics Subject Classification.* Primary: 35L05, 35B40; Secondary: 35L70, 35B35 .

Key words and phrases. Wave equation, linear damping, Decay rate.

This work is supported by the Tunisian Ministry for Scientific Research and Technology within the LAB-STI 02 program.

with localized dissipations. For the Klein-Gordon equations the energy functional it self contains the L^2 norm and boundedness of L^2 norm of the solution is trivial. Thus under some assumptions on the support of the damper a we can show that the energy decays exponentially while for the system (1.1) the energy decay rate is weaker and more delicate.

In the case when $a(x) \geq \epsilon_0 > 0$ in all of Ω we know that

$$E_u(t) \leq C_0 (1+t)^{-1} \text{ and } \|u(t)\|_{L^2}^2 \leq C_0, \text{ for all } t \geq 0, \tag{1.4}$$

for weak solution u to the system (1.1) with initial data in $H_0^1(\Omega) \times L^2(\Omega)$.

Nakao in [26] obtained the same estimates in (1.4) for a damper a which is positive near some part of the boundary (Lions's condition) and near infinity.

On the other hand, Dan-Shibata [8] studied the local energy decay estimates for the compactly supported weak solutions of (1.1) with $a(x) = 1$

$$\int_{\Omega \cap B_R} \left(|\nabla u(t,x)|^2 + |\partial_t u(t,x)|^2 \right) dx \leq C (1+t)^{-d}, \tag{1.5}$$

where $B_R = \left\{ x \in \mathbb{R}^d, |x| < R \right\}$. For damping localized in a compact set of Ω: Nakao in [25] proved that the local energy decays exponentially if d is odd and polynomially if d is even under the Lions's geometric condition. Combining the definition of a non-trapping obstacle and the geometric control condition of Bardos–Lebeau–Rauch [5], Aloui and Khenissi [2] introduced the exterior geometric control condition which is a sufficient condition for the stabilization of the local energy, more precisely they proved that the local energy decays exponentially if the space dimension d is odd. In [21] Khenissi showed that the estimate (1.5) holds if d is even. Recently in [9], using a nonlinear internal localized damping, Daoulatli obtained various decay rates, depending on the behavior of the damping term. We also mention the result of Daoulatli-Dehman-Khenissi [10] on the behaviors of the local energy for solutions of the Lamé system in exterior domain. Finally, we quote the result of Bchatnia and Daoulatli [4] on the decay rates of the local energy for solutions of the wave equation with localized time dependent damping in exterior domain.

Furthermore Ikehata and Matsuyama in [18] obtained a more precise decay estimate for the total energy of solutions of the problem (1.1) with $a(x) = 1$ and for weighted initial data

$$E_u(t) \leq C_2 (1+t)^{-2} I_1 \text{ and } \|u(t)\|_{L^2}^2 \leq C_2 (1+t)^{-1} I_1 \text{ for all } t \geq 0. \tag{1.6}$$

Especially this estimate seems sharp for $d = 2$ as compared with that of [8].

Ikehata in [17] derived a fast decay rate like (1.6) for solutions of the system (1.1) with weighted initial data and assuming that $a(x) \geq \epsilon_0 > 0$ at infinity and $O = \mathbb{R}^d \backslash \Omega$ is star shaped with respect to the origin.

More recently in [12] Daoulatli prove that for a damper a positive near infinity and under the geometric control condition of Bardos et al [5], the estimates in (1.4) hold for all solutions of the system (1.1) with initial data in $H_0^1(\Omega) \times L^2(\Omega)$ and show that the estimates in (1.6) hold for all solutions of the system (1.1) with weighted initial data. Moreover he prove that for every $p \in \mathbb{N}^*$ there exists a initial data in $H_0^1(\Omega) \times L^2(\Omega)$ such that the solution v of (1.1) verifies

$$E_v(t) \leq C (1+t)^{-p} \text{ and } \|v(t)\|_{L^2}^2 \leq C (1+t)^{-p+1}, \text{ for all } t \geq 0$$

and for some $C > 0$ depending on the initial data.

Now we give a summary of results on the asymptotic behavior of the energy of solutions of the nonlinear system (1.1) in the free space $\mathbb{R}^d$ and for a globally distributed damping.

For the Klein Gordon equation a polynomial decay rate is derived by Nakao [28] for compactly supported initial data and by Mochizuki and Motai [24] for weighted initial data. More precisely they show that if $1 < r < 1 + \frac{2}{d}$ the energy decay according to

$$E_u(t) \leq C(1+t)^{-\gamma}$$

where $\gamma = \frac{2+d-dr}{r-1}$. If $r > 1 + \frac{2}{d}$, Mochizuki and Motai [24] establishes a complementary non-decay result for a dense set of initial data in $H^1(\mathbb{R}^d) \times L^2(\mathbb{R}^d)$.

For the wave equation first we quote the result of Ono [31], Ono in [31] considered the wave equation with the following damping $\partial_t u + g(\partial_t u)$ with g superlinear and has a polynomial growth. He showed the polynomial decay of the energy. We note that in this case the L^2 norm of the time derivative on $\mathbb{R}_+ \times \mathbb{R}^d$ of the solution is bounded by the energy of the initial data. Mochizuki and Motai in [24] obtained a logarithmic decay rate when $1 < r \leq 1 + \frac{2}{d}$ and for a kind of weighted initial data. The corresponding non-decay result in [24] requires $r > 1 + \frac{2}{d-1}$. Todorova and Yordanov in [35] showed that for compactly supported initial data there exists a positive constant τ such that

$$E_u(t) \leq C(1+t)^{-\tau}$$

when $1 < r \leq 1 + \frac{2}{d+1}$ and $d \geq 3$.

In the case of exterior domain we mention the result of Nakao and Jung [30] which consider a dissipation which is allowed to be nonlinear only in a ball, but outside that ball the dissipation must be linear. For the generalized Klein Gordon equation we quote the result of Nakao [29].

For solutions of the system (1.1) and Ω is a exterior domain no results seems to be known.

For another type of total energy decay property we refer the reader to [16, 20, 33, 3, 32, 15] and references therein.

Before introducing our results we shall state several assumptions:

Hyp A: There exists $L > 0$ such that

$$a(x) > \epsilon_0 > 0 \text{ for } |x| \geq L.$$

Definition 1. *(ω, T) geometrically controls Ω, i.e. every generalized geodesic travelling with speed 1 and issued at $t = 0$, enters the set ω in a time $t < T$.*

This condition is called Geometric Control Condition (see e.g.[5]). We shall relate the open subset ω with the damper a by

$$\omega = \{x \in \Omega; a(x) > \epsilon_0 > 0\}.$$

We note that according to [5] and [7] the Geometric Control Condition of Bardos et al is a necessary and sufficient condition for the stabilization of the wave equation in bounded domain.

In this paper we assume that the initial data (u_0, u_1) in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ and we prove that for a damper a positive at infinity and under the geometric control condition of Bardos et al [5]:

(1) If $1 < r \leq 1 + \frac{2}{d}$ and

$$\left\| (\ln(b+q))^{\frac{\gamma}{2}} \nabla u_0 \right\|_{L^2}^2 + \left\| (\ln(b+q))^{\frac{\gamma}{2}} u_1 \right\|_{L^2}^2 < +\infty,$$

then the energy decays as

$$E_u(t) \leq C_0 (\ln(b+t))^{-\gamma}$$

(2) If $1 < r < 1 + \frac{2}{d}$ and

$$\left\| (1+q)^{\frac{\gamma}{2}} \nabla u_0 \right\|_{L^2}^2 + \left\| (1+q)^{\frac{\gamma}{2}} u_1 \right\|_{L^2}^2 < +\infty$$

then the energy decays as

$$E_u(t) \leq C_1 (1+t)^{-\gamma}$$

(3) If $1 < r < 1 + \frac{2}{d}$ and the initial data has a compact support, then the enegy decays as

$$E_u(t) \leq C_1 (1+t)^{-\gamma_1}$$

Throughout this paper we use the following notations

$$q(x) = \left(1 + |x|^2\right)^{\frac{1}{2}}, \text{ for } x \in \Omega$$

Let $\psi \in C_0^\infty\left(\mathbb{R}^d\right)$ such that $0 \leq \psi \leq 1$ and

$$\psi(x) = \begin{cases} 1 & \text{for } |x| \leq L \\ 0 & \text{for } |x| \geq 2L \end{cases}.$$

We set

$$p = \begin{cases} 2(r+1) & \text{if } d \leq 3 \\ \frac{2d}{d-2} & \text{if } d \geq 4 \end{cases}$$

Now we state the results of this paper.

Theorem 1. *We assume that Hyp A holds and* (ω, T) *geometrically controls* Ω. *Let*

$$\begin{array}{ll} \gamma > 0 & \textit{if } 1 < r < 1 + \frac{2}{d} \\ 0 < \gamma < \frac{2}{r-1} & \textit{if } r = 1 + \frac{2}{d} \end{array}$$

and

$$\ln b = \max\left((2(r+1))^{r+1}, \gamma - 1, \frac{\gamma - 1 + \sqrt{|(\gamma-1)(\gamma+3)|}}{2}, \frac{\gamma - r}{r-1}, \left(\frac{4(r+1)\gamma}{1-\delta_0}\right)^{r+1} \right)$$

where $0 < \delta_0 < 1$. *Then there exists* $C_0 > 0$ *such that the following estimate*

$$E_u(t) \leq C_0 (\ln(b+t))^{-\gamma} I_0, \ \textit{for all } t \geq 0$$

holds for every solution u *of* (1.1) *with initial data* (u_0, u_1) *in* $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$, *such that*

$$\left\| (\ln(b+q))^{\frac{\gamma}{2}} \nabla u_0 \right\|_{L^2}^2 + \left\| (\ln(b+q))^{\frac{\gamma}{2}} u_1 \right\|_{L^2}^2 < +\infty,$$

where

$$I_0 = \|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r} + \|u_0\|_{L^{r+1}}^{r+1} + \left\| (\ln(b+q))^{\frac{\gamma}{2}} \nabla u_0 \right\|_{L^2}^2$$
$$+ \left\| (\ln(b+q))^{\frac{\gamma}{2}} u_1 \right\|_{L^2}^2 + \left(\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r} \right)^{\frac{p}{2}} + 1$$

Moreover we have

$$\begin{aligned}
&\int_{\Omega}\left(\ln\left(b+q\left(x\right)+t\right)\right)^{\gamma}\left(\left|\nabla u\right|^{2}+\left|\partial_{t}u\right|^{2}\right)dx\leq C_{0}I_{0},\\
&\int_{0}^{+\infty}\int_{\Omega}\tfrac{\left(\ln(b+q(x)+s)\right)^{\gamma-1}}{b+q(x)+s}\left(\left|\nabla u\right|^{2}+\left|\partial_{t}u\right|^{2}\right)dxds\leq C_{0}I_{0},\\
&\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\left(\ln\left(b+q\left(x\right)+t\right)\right)^{\gamma}\left|\partial_{t}u\right|^{r+1}dxds\leq C_{0}I_{0},\\
&\int_{\Omega}a\left(x\right)\tfrac{\left(\ln(b+q(x)+t)\right)^{\gamma-1}}{(b+q(x)+t)^{2}}\left|u\left(t\right)\right|^{2}dx\leq C_{0}I_{0},\\
&\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\tfrac{\left(\ln(b+q(x)+t)\right)^{\gamma-1}}{(b+q(x)+t)^{3}}\left|u\left(t\right)\right|^{2}dxds\leq C_{0}I_{0},\\
&\int_{\Omega}a\left(x\right)\tfrac{\left(\ln(b+q(x)+t)\right)^{\gamma-r}}{(b+q(x)+t)^{r}}\left|u\left(t\right)\right|^{r+1}dx\leq C_{0}I_{0},\\
&\textit{and }\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\tfrac{\left(\ln(b+q(x)+t)\right)^{\gamma-r}}{(b+q(x)+t)^{r+1}}\left|u\left(t\right)\right|^{r+1}dxds\leq C_{0}I_{0}.
\end{aligned}$$

Theorem 2. *We assume that Hyp A holds and (ω,T) geometrically controls Ω. We suppose that $1<r<1+\frac{2}{d}$. Let $0<\delta_0\ll\frac{1}{2}$ and $0<\gamma<\min\left(\frac{\frac{1}{2}-\delta_0}{k},\frac{d+2-dr}{r-1},\frac{p-2r}{r-1}\right)$ with*

$$\begin{aligned}
k=&\tfrac{5(1+\delta_0)r^2+8\left(\frac{1}{2}-\delta_0\right)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)}{10r(r+1)}\\
&+\tfrac{\sqrt{\left(5(1+\delta_0)r^2+8\left(\frac{1}{2}-\delta_0\right)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)^2-160\left((1+\delta_0)\left(\frac{1}{2}-\delta_0\right)\right)r^2(r+1)}}{10r(r+1)}
\end{aligned}$$

Then there exists $C_1>0$ such that the following estimate

$$E_u\left(t\right)\leq C_1\left(1+t\right)^{-\gamma}I_1,\ \textit{for all } t\geq 0$$

holds for every solution u of (1.1) with initial data (u_0,u_1) in $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$, such that

$$\left\|\left(1+q\right)^{\frac{\gamma}{2}}\nabla u_0\right\|_{L^2}^2+\left\|\left(1+q\right)^{\frac{\gamma}{2}}u_1\right\|_{L^2}^2<+\infty$$

where

$$\begin{aligned}
I_1=&\left\|u_0\right\|_{H^2}^2+\left\|u_1\right\|_{H^1}^2+\left\|u_1\right\|_{H^1}^{2r}+\left\|u_0\right\|_{L^{r+1}}^{r+1}+\left\|\left(1+q\right)^{\frac{\gamma}{2}}\nabla u_0\right\|_{L^2}^2\\
&+\left\|\left(1+q\right)^{\frac{\gamma}{2}}u_1\right\|_{L^2}^2+\left(\left\|u_0\right\|_{H^2}^2+\left\|u_1\right\|_{H^1}^2+\left\|u_1\right\|_{H^1}^{2r}\right)^{\frac{p}{2}}+1
\end{aligned}$$

Moreover we have

$$\begin{aligned}
&\int_{\Omega}\left(1+q\left(x\right)+t\right)^{\gamma}\left(\left|\nabla u\right|^{2}+\left|\partial_{t}u\right|^{2}\right)dx\leq C_{1}I_{1},\\
&\int_{0}^{+\infty}\int_{\Omega}\left(1+q\left(x\right)+s\right)^{\gamma-1}\left(\left|\nabla u\right|^{2}+\left|\partial_{t}u\right|^{2}\right)dxds\leq C_{1}I_{1},\\
&\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\left(1+q\left(x\right)+s\right)^{\gamma}\left|\partial_{t}u\right|^{r+1}dxds\leq C_{1}I_{1},\\
&\int_{\Omega}a\left(x\right)\left(1+q\left(x\right)+t\right)^{\gamma-2}\left|u\left(t\right)\right|^{2}dx\leq C_{1}I_{1},\\
&\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\left(1+q\left(x\right)+s\right)^{\gamma-3}\left|u\left(t\right)\right|^{2}dxds\leq C_{1}I_{1},\\
&\int_{\Omega}a\left(x\right)\left(1+q\left(x\right)+t\right)^{\gamma-r}\left|u\left(t\right)\right|^{r+1}dx\leq C_{1}I_{1},\\
&\textit{and }\int_{0}^{+\infty}\int_{\Omega}a\left(x\right)\left(1+q\left(x\right)+s\right)^{\gamma-r-1}\left|u\left(t\right)\right|^{r+1}dxds\leq C_{1}I_{1}.
\end{aligned}$$

Theorem 3. *We assume that Hyp A holds and* (ω,T) *geometrically controls* Ω*. Let* $0<\delta_0\ll 1$ *and* $R\geq 1$ *such that* $O\subset B_R$*. Then there exists* $C_2>0$ *such that the following estimate*

$$E_u(t)\leq C_2\left(\frac{R}{R+t}\right)^{\gamma} I_2,\ \textit{for all } t\geq 0$$

holds for every solution u *of* (1.1) *with initial data* (u_0,u_1) *in* $H_0^1(\Omega)\cap H^2(\Omega)\times H_0^1(\Omega)$*, such that support of* (u_0,u_1) *is contained in* B_R*, where*

$$I_2=\|u_0\|_{H^2}^2+\|u_1\|_{H^1}^2+\|u_1\|_{H^1}^{2r}+\|u_0\|_{L^{r+1}}^{r+1}+\left(\|u_0\|_{H^2}^2+\|u_1\|_{H^1}^2+\|u_1\|_{H^1}^{2r}\right)^{\frac{p}{2}}+1$$

and $0<\gamma<\min\left(\frac{\frac{1}{2}-\delta_0}{k},\frac{d+2-dr}{r-1},\frac{p-2r}{r-1}\right)$ *with*

$$k=\tfrac{5(1+\delta_0)r^2+8(1-\delta_0)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)}{10r(r+1)}$$
$$+\tfrac{\sqrt{\left(5(1+\delta_0)r^2+8(1-\delta_0)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)^2-160r^2(r+1)\left(1-\delta_0^2\right)}}{10r(r+1)}$$

Moreover we have

$$\begin{aligned}
&\textstyle\int_0^{+\infty}(R+t)^{\gamma-1}E_u(t)\,dt\leq C_2R^{\gamma}I_2,\\
&\textstyle\int_0^{+\infty}(R+t)^{\gamma}\int_\Omega a(x)\left|\partial_t u\right|^{r+1}dxdt\leq C_2R^{\gamma}I_2,\\
&\textstyle\int_\Omega a(x)\left|u(t)\right|^2dx\leq C_2R^{\gamma}(R+t)^{2-\gamma}I_2,\\
&\textstyle\int_\Omega a(x)\left|u(t)\right|^{r+1}dx\leq C_2R^{\gamma}(R+t)^{r-\gamma}I_2,\\
&\textstyle\int_0^{+\infty}(R+t)^{\gamma-3}\int_\Omega a(x)\left|u(t)\right|^2dxdt\leq C_2R^{\gamma}I_2,\\
&\textit{and }\textstyle\int_0^{+\infty}(R+t)^{\gamma-r-1}\int_\Omega a(x)\left|u(t)\right|^{r+1}dxdt\leq C_2R^{\gamma}I_2.
\end{aligned}$$

First we remark that the constant C_2 depends on R, but we can prove the same result with constant independent of R. We note that that the rate of decay in the case of initial data with compact support is better or equal to the one obtained for weighted initial data with equivalent polynomial weight.

The key idea is to use the following auxiliary functional

$$\begin{aligned}
X(t)&=\textstyle\int_\Omega f_0(\alpha q(x)+t)\,v(t)\,\partial_t v(t)\,dx+k_1\int_\Omega f_1(\alpha q(x)+t)\,a(x)\left|u(t)\right|^2dx\\
&\quad+k_2\textstyle\int_\Omega a(x)f_2(\alpha q(x)+t)\left|u(t)\right|^{r+1}dx+k\int_\Omega f(\alpha q(x)+t)\left(\left|\nabla u(t)\right|^2+\left|\partial_t u(t)\right|^2\right)dx,
\end{aligned}$$

$\alpha\geq 0$, u is the solution of (1.1) with initial data in $H_0^1(\Omega)\times L^2(\Omega)$ and $v=(1-\psi)\,u$ where k_1,k_2 and k are positive constants and

$$f_0\simeq f',\ \ f_1\simeq -f''\text{ and }f_2\simeq\left(f'\right)^r(f)^{-r+1}$$

where f is some positive, concave function. Then to show a weighted observability estimates for the local energy using technics based on the notion of microlocal defect measure and exploiting their properties. Especially the fact that the measure associated to a bounded sequence of solutions of the conservative wave equation propagates along the generalized bicharacteristic flow of Melrose-Sjöstrand.

We note that if we take initial data with finite weighted energy with weight slower than $\ln(b+q)$, for example $\ln\ln(b+q)$, then the energy decays like $(\ln\ln(b+t))^{-1}$.

The rest of the paper is organized as follows. In section 2 we present some results on the weighted energy and we give a weighted observability estimate for the local energy. The section 3 is devoted to the proof of theorem 1and in section 4 we give the proof of theorem 2, finally in the section 5 we give the proof of theorem 3.

2. Weighted observability estimate

The next result concern the weighted energy estimate for solutions of (1.1) with weighted initial data.

Proposition 1. *Let φ be a positive function in $C^2(\mathbb{R}_+)$ such that $\varphi' \in L^\infty(\mathbb{R}_+)$ and $\varphi'' \in L^\infty(\mathbb{R}_+)$. Let u be a solution of* (1.1) *with initial data (u_0, u_1) in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ such that*

$$\int_\Omega \varphi(\mu q(x)) \left(|\nabla u_0|^2 + |u_1|^2\right) dx < \infty.$$

Setting

$$E_\varphi(u)(t) = \frac{1}{2}\int_\Omega \varphi(\mu q(x) + \lambda t) \left(|\nabla u|^2 + |\partial_t u|^2\right) dx,$$

where $\lambda, \mu \geq 0$. Then

$$\sqrt{\varphi}\nabla u \in L^\infty_{loc}\left(\mathbb{R}_+, \left(L^2(\Omega)\right)^d\right) \text{ and } \sqrt{\varphi}\partial_t u \in L^\infty_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right) \tag{2.1}$$

Moreover we have

$$\begin{array}{l} E_\varphi(u)(t+T) + \int_t^{t+T}\int_\Omega a(x)\varphi\,|\partial_t u|^{r+1}\,dxds \\ \leq E_\varphi(u)(t) + \frac{\lambda+\mu}{2}\int_t^{t+T}\int_\Omega |\varphi'| \left(|\nabla u(s)|^2 + |\partial_t u(s)|^2\right) dxds. \end{array} \tag{2.2}$$

for every $t \geq 0$ and $T \geq 0$.

Proof. Let $n \in \mathbb{N}^*$. We define

$$g_n(s) = g \circ \left(I + n^{-1}g\right)^{-1}(s) = n\left(s - \left(I + n^{-1}g\right)^{-1}(s)\right)$$

the Yosida approximation of $g : s \longmapsto |s|^{r-1} s$. g_n is monotone increasing and globally Lipschitz. Let w be the solution of the following system

$$\left\{\begin{array}{ll} \partial_t^2 w - \Delta w + a(x)(1+\varphi)^{\frac{1}{2}} g_n\left((1+\varphi)^{-\frac{1}{2}}\partial_t w + h(t,x)\right) = f(t,x) & \mathbb{R}_+ \times \Omega \\ w = 0 & \mathbb{R}_+ \times \partial\Omega \\ (w(0), \partial_t w(0)) = (w_0, w_1) \in H_D(\Omega) \times L^2(\Omega) & \end{array}\right. \tag{2.3}$$

with $f \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right)$ and $(1+\varphi)^{\frac{1}{2}} h \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right)$, where $H_D(\Omega)$ the completion of $C_0^\infty(\Omega)$ with respect to the norme

$$\|\varphi_0\|^2_{H_D} = \int_\Omega |\nabla \varphi_0|^2\, dx.$$

g_n is a global Lipschitz function, therefore

$$a(x)(1+\varphi)^{\frac{1}{2}} g_n\left((1+\varphi)^{-\frac{1}{2}}\partial_t w + h(t,x)\right) \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right).$$

Using the fact that the function $f \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right)$, we infer that the unique solution of (2.3)

$$w \in C(\mathbb{R}_+, H_D(\Omega)) \text{ and } \partial_t w \in C\left(\mathbb{R}_+, L^2(\Omega)\right),$$

and the following energy identity

$$\begin{aligned} & E_w(t) + \int_0^t \int_\Omega a(x)(1+\varphi)^{\frac{1}{2}} g_n\left((1+\varphi)^{-\frac{1}{2}}\partial_t w + h(s,x)\right)\partial_t w\,dx\,ds \\ = \; & E_w(0) + \int_0^t \int_\Omega f(t,x)\,\partial_t w\,dx\,ds \end{aligned} \tag{2.4}$$

holds for every $t \geq 0$.

Let u be the solution of the following system

$$\left\{\begin{array}{ll} \partial_t^2 u - \Delta u + a(x)\, g_n(\partial_t u) = 0 & \mathbb{R}_+ \times \Omega \\ u = 0 & \mathbb{R}_+ \times \partial\Omega \\ (u(0), \partial_t u(0)) = (u_0, u_1) & \end{array}\right. \tag{2.5}$$

with (u_0, u_1) in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ such that

$$\int_\Omega \varphi(\mu q(x))\left(|\nabla u_0|^2 + |u_1|^2\right)dx < \infty.$$

The function g_n is globally Lipschitz, hence

$$u \in L^\infty\left([0,T], H_0^1(\Omega) \cap H^2(\Omega)\right) \cap W^{1,\infty}\left([0,T], H_0^1(\Omega)\right),$$

moreover we have the following energy identity

$$E_u(t) + \int_0^t \int_\Omega a(x)\, g_n(\partial_t u)\,\partial_t u\,dx\,ds = E_u(0) \tag{2.6}$$

We set $v = (1+\varphi)^{\frac{1}{2}} u$. Therefore v satisfies

$$\left\{\begin{array}{ll} \partial_t^2 v - \Delta v + a(x)(1+\varphi)^{\frac{1}{2}} g_n\left((1+\varphi)^{-\frac{1}{2}}\partial_t v + h(t,x)\right) = f(t,x) & \mathbb{R}_+ \times \Omega \\ v = 0 & \mathbb{R}_+ \times \partial\Omega \\ (v(0), \partial_t v(0)) = (v_0, v_1) & \end{array}\right. \tag{2.7}$$

with

$$\begin{aligned} (v_0, v_1) &= \left((1+\varphi)^{\frac{1}{2}} u_0, \frac{\lambda}{2}(1+\varphi)^{-\frac{1}{2}}\varphi' u_0 + (1+\varphi)^{\frac{1}{2}} u_1\right) \\ h(t,x) &= -\frac{\lambda}{2}(1+\varphi)^{-1}\varphi' u \end{aligned}$$

and

$$\begin{aligned} f(t,x) = {} & \tfrac{\mu}{2}\left[\mu\left(\varphi''(1+\varphi)^{-\frac{1}{2}} - \tfrac{1}{2}(\varphi')^2(1+\varphi)^{-3/2}\right)\tfrac{|x|^2}{q^2} + \left(\tfrac{d}{q} - \tfrac{|x|^2}{q^3}\right)\varphi'(1+\varphi)^{-\frac{1}{2}}\right]u \\ & + \tfrac{\lambda^2}{2}\left[\varphi''(1+\varphi)^{-\frac{1}{2}} - \tfrac{1}{2}(\varphi')^2(1+\varphi)^{-3/2}\right]u + \varphi'(1+\varphi)^{-\frac{1}{2}}\left(\lambda\partial_t u + \mu\tfrac{x\cdot\nabla u}{q}\right) \end{aligned}$$

It is easy to see that

$$\begin{array}{l} (v_0, v_1) \in H_D(\Omega) \times L^2(\Omega) \\ (1+\varphi)^{\frac{1}{2}} h(t,x) \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right) \\ f(t,x) \in L^2_{loc}\left(\mathbb{R}_+, L^2(\Omega)\right) \end{array}$$

Therefore using (2.4), we deduce that

$$\begin{aligned} & E_v(t) + \textstyle\int_0^t \int_\Omega a(x)(1+\varphi)\, g_n(\partial_t u)\,\partial_t u\,dx\,ds \\ & = E_v(0) + \textstyle\int_0^t \int_\Omega (1+\varphi)^{\frac{1}{2}} f(t,x)\,\partial_t u\,dx\,ds + \frac{\lambda}{2}\int_0^t \int_\Omega (1+\varphi)^{-\frac{1}{2}} f(t,x)\,\varphi' u\,dx\,ds \\ & -\tfrac{\lambda}{2}\textstyle\int_0^t \int_\Omega a(x)\,\varphi' g_n(\partial_t u)\,u\,dx\,ds \end{aligned} \tag{2.8}$$

It clear that there exists a positive constant $C = C(\lambda, \mu, \varphi)$ such that

$$\left|\int_0^t \int_\Omega (1+\varphi)^{\frac{1}{2}} f(t,x)\, \partial_t u dxds\right| \leq C \int_0^t \int_\Omega |u|^2 + |\partial_t u|^2 + |\nabla u|^2 dxds,$$

$$\left|\frac{\lambda}{2} \int_0^t \int_\Omega (1+\varphi)^{-\frac{1}{2}} f(t,x)\, \varphi' u dxds\right| \leq C \int_0^t \int_\Omega |u|^2 + |\partial_t u|^2 + |\nabla u|^2 dxds$$

and

$$\left|\frac{\lambda}{2} \int_0^t \int_\Omega a(x)\, \varphi' g_n(\partial_t u)\, u dxds\right| \leq C \int_0^t \int_\Omega a(x)\, |u|^{r+1} + a(x) \left|\left(I + n^{-1} g\right)^{-1} (\partial_t u)\right|^{r+1} dxds$$

Now using (2.6) and the fact that

$$g_n(\partial_t u)\, \partial_t u \geq \left|\left(I + n^{-1} g\right)^{-1} (\partial_t u)\right|^{r+1}$$

we infer that

$$\int_0^t \int_\Omega a(x) \left|\left(I + n^{-1} g\right)^{-1} (\partial_t u)\right|^{r+1} dxds \leq E_u(0)$$

and

$$\int_0^t \int_\Omega |\partial_t u|^2 + |\nabla u|^2 dxds \leq (1+t)\, E_u(0)$$

We have

$$\begin{gathered} \int_\Omega |u(s)|^2 dx \leq C(1+s)^2 \left(E_u(0) + \|u_0\|_{L^2}^2\right) \\ \text{and} \\ \int_\Omega a(x)\, |u(s)|^{r+1} dx \leq C(1+s)^\nu \left(E_u(0) + \|u_0\|_{L^2}^2\right)^{\frac{r+1}{2}} \end{gathered} \tag{2.9}$$

with $\nu = \frac{2(p-r-1)}{(r+1)(p-2)}$ and

$$p = \begin{cases} 2(r+1) & \text{if } d \leq 2 \\ \frac{2d}{d-2} & \text{if } d \geq 3 \end{cases}$$

Therefore

$$\begin{gathered} \int_0^t \int_\Omega |u|^2 dxds \leq C(1+t)^3 \left(E_u(0) + \|u_0\|_{L^2}^2\right) \\ \text{and} \\ \int_0^t \int_\Omega a(x)\, |u|^{r+1} dxds \leq C(1+t)^{\nu+1} \left(E_u(0) + \|u_0\|_{L^2}^2\right)^{\frac{r+1}{2}} \end{gathered}$$

Combining the estimates above with (2.8), we obtain

$$\begin{aligned} & E_v(t) + \int_0^t \int_\Omega a(x)(1+\varphi)\, g_n(\partial_t u)\, \partial_t u dxds \\ & \leq C(1+t)^\alpha \left(E_v(0) + E_u(0) + \left(E_u(0) + \|u_0\|_{L^2}^2\right)^{\frac{r+1}{2}} + \|u_0\|_{L^2}^2\right) \end{aligned} \tag{2.10}$$

with $\alpha = \max(3, \nu+1)$. It is clear that

$$E_\varphi(u)(t) \leq 2E_v(t) + C\, \|u(t)\|_{L^2}^2$$

Therefore combining the estimate above with (2.10) and (2.9) we obtain

$$\begin{aligned} & E_\varphi(u)(t) + \int_0^t \int_\Omega a(x)(1+\varphi)\, g_n(\partial_t u)\, \partial_t u dxds \\ & \leq C(1+t)^\alpha \left(E_\varphi(u)(0) + E_u(0) + \left(E_u(0) + \|u_0\|_{L^2}^2\right)^{\frac{r+1}{2}} + \|u_0\|_{L^2}^2\right) \end{aligned} \tag{2.11}$$

We note that $C = C(\lambda, \mu, \varphi)$.

Let (u_0, u_1) in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ such that

$$\int_\Omega \varphi(\mu q(x)) \left(|\nabla u_0|^2 + |u_1|^2\right) dx < \infty$$

and consider the system

$$\begin{cases} \partial_t^2 u_n - \Delta u_n + a(x) g_n(\partial_t u_n) = 0 & \mathbb{R}_+ \times \Omega \\ u_n = 0 & \mathbb{R}_+ \times \partial\Omega \\ (u_n(0), \partial_t u_k(0)) = (u_0, u_1) & \end{cases} \tag{2.12}$$

From (1.2) and (1.3), we infer that there exists u such that

$$\begin{array}{l} u_n \underset{n\to+\infty}{\longrightarrow} u \text{ in the weak star topology of } L^\infty\left([0,T], H_0^1(\Omega) \cap H^2(\Omega)\right) \\ \partial_t u_n \underset{n\to+\infty}{\longrightarrow} \partial_t u \text{ in the weak star topology of } L^\infty\left([0,T], H_0^1(\Omega)\right) \\ \left(I + n^{-1} g\right)^{-1}(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} \partial_t u \text{ in the weak topology of } L^{r+1}((0,T) \times \Omega, a) \\ g_n(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} \psi \text{ in the weak topology of } L^{\frac{r+1}{r}}((0,T) \times \Omega, a) \end{array} \tag{2.13}$$

To show that, $\psi = g(\partial_t u)$, we proceed as in [23, P55-56]. By a classical compactness argument, we can show that there exists a subsequence of (u_n) still denoted by (u_n), such that

$$\partial_t u_n \underset{n\to+\infty}{\longrightarrow} \partial_t u \text{ strongly in } L^2(K). \tag{2.14}$$

for a given compact subset K of $(0,T) \times \Omega$. Therefore we can assume that

$$\partial_t u_n \underset{n\to+\infty}{\longrightarrow} \partial_t u, \text{ a.e. in } K.$$

Since the function $s \longmapsto \left(I + n^{-1} g\right)^{-1}(s)$, is non-expansive on $\mathbb{R}$, we obtain

$$\left(I + n^{-1} g\right)^{-1}(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} \partial_t u, \text{ a.e. in } K.$$

Hence

$$g_n(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} g(\partial_t u), \text{ a.e. in } K.$$

This enough to gives $\psi = g(\partial_t u)$. Therefore u is a solution of (1.1) with initial data in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ such that

$$\int_\Omega \varphi(\mu q(x)) \left(|\nabla u_0|^2 + |u_1|^2\right) dx < \infty.$$

Now using (2.11) and the fact that

$$g_n(\partial_t u_n) \partial_t u_n \geq \left|\left(I + n^{-1} g\right)^{-1}(\partial_t u_n)\right|^{r+1}$$

we infer that

$$\begin{array}{l} \sqrt{1+\varphi}\,\partial_{x_i} u_n \underset{n\to+\infty}{\longrightarrow} \psi_i \text{ in the weak star topology of } L^\infty\left([0,T], L^2(\Omega)\right),\ i \in \{1,..,d\} \\ \sqrt{1+\varphi}\,\partial_t u_n \underset{n\to+\infty}{\longrightarrow} \phi_1 \text{ in the weak star topology of } L^\infty\left([0,T], L^2(\Omega)\right) \\ (a(1+\varphi))^{\frac{1}{r+1}} \left(I + n^{-1} g\right)^{-1}(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} \phi_2 \text{ in the weak topology of } L^{r+1}((0,T) \times \Omega) \end{array}$$

Now we show that

$$\psi_i = \sqrt{1+\varphi}\,\partial_{x_i} u,\ \phi_1 = \sqrt{1+\varphi}\,\partial_t u \text{ and } \phi_2 = (a(1+\varphi))^{\frac{1}{r+1}} \partial_t u$$

Let K be a compact set of $(0,T)\times\Omega$. Using (2.14) and extracting a subsequence if necessary

$$\partial_t u_n \underset{n\to+\infty}{\longrightarrow} \partial_t u, \text{ a.e. in } K,$$

hence

$$\sqrt{1+\varphi}\partial_t u_n \underset{n\to+\infty}{\longrightarrow} \sqrt{1+\varphi}\partial_t u, \text{ a.e. in } K,$$

and using the fact that the function $s \longmapsto \left(I+n^{-1}g\right)^{-1}(s)$, is non-expansive on $\mathbb{R}$, we obtain

$$(a(1+\varphi))^{\frac{1}{r+1}}\left(I+n^{-1}g\right)^{-1}(\partial_t u_n) \underset{n\to+\infty}{\longrightarrow} (a(1+\varphi))^{\frac{1}{r+1}}\partial_t u, \text{ a.e. in } K,$$

This is enough to imply

$$\phi_1=\sqrt{1+\varphi}\partial_t u \text{ and } \phi_2=(a(1+\varphi))^{\frac{1}{r+1}}\partial_t u$$

From (2.13) and by a classical compactness argument, we can show that there exists a subsequence of (u_n) still denoted by (u_n), such that

$$\partial_{x_i} u_n \underset{n\to+\infty}{\longrightarrow} \partial_{x_i} u \text{ strongly in } L^2(K).$$

Therefore extracting a subsequence if necessary

$$\partial_{x_i} u_n \underset{n\to+\infty}{\longrightarrow} \partial_{x_i} u, \text{ a.e. in } K,$$

which gives

$$\sqrt{1+\varphi}\partial_{x_i} u_n \underset{n\to+\infty}{\longrightarrow} \sqrt{1+\varphi}\partial_{x_i} u\ , \text{ a.e. in } K.$$

We conclude that

$$\psi_i=\sqrt{1+\varphi}\partial_{x_i} u,\ i\in\{1,..,d\}.$$

Therefore

$$\sqrt{\varphi}\nabla u\in L^\infty_{loc}\left(\mathbb{R}_+,\left(L^2(\Omega)\right)^d\right) \text{ and } \sqrt{\varphi}\partial_t u\in L^\infty_{loc}\left(\mathbb{R}_+,L^2(\Omega)\right) \tag{2.15}$$

We remind that

$$u\in L^\infty_{loc}\left(\mathbb{R}_+,H^1_0(\Omega)\cap H^2(\Omega)\right)\cap W^{1,\infty}\left(\mathbb{R}_+,H^1_0(\Omega)\right)\cap W^{2,\infty}\left(\mathbb{R}_+,L^2(\Omega)\right).$$

Let $R\gg 1$ and setting $S(R)=\partial B_R$. It is clear that

$$\begin{aligned}
&\tfrac{1}{2}\tfrac{d}{dt}\textstyle\int_{\Omega\cap B_R}\varphi\left(|\nabla u(t)|^2+|\partial_t u(t)|^2\right)dx+\int_{\Omega\cap B_R}a(x)\varphi|\partial_t u(t)|^{r+1}dx\\
&=\tfrac{\lambda}{2}\textstyle\int_{\Omega\cap B_R}\varphi'\left(|\nabla u(t)|^2+|\partial_t u(t)|^2\right)dx+\int_{\Omega\cap B_R}\varphi\nabla u(t)\cdot\nabla\partial_t u(t)+\varphi\partial_t u(t)\,\partial_t^2 u(t)\,dx\\
&=\tfrac{\lambda}{2}\textstyle\int_{\Omega\cap B_R}\varphi'\left(|\nabla u(t)|^2+|\partial_t u(t)|^2\right)dx-\mu\int_{\Omega\cap B_R}\varphi'\tfrac{x\cdot\nabla u(t)}{q(x)}\partial_t u(t)\,dx+\int_{S(R)}\varphi\tfrac{x\cdot\nabla u(t)}{R}\partial_t u(t)\,dS,
\end{aligned}$$

Integrating the estimate above between t and $t+T$, we obtain

$$\begin{aligned}
&\textstyle\int_{\Omega\cap B_R}\varphi\left(|\nabla u(t+T)|^2+|\partial_t u(t+T)|^2\right)dx+\int_t^{t+T}\int_{\Omega\cap B_R}a(x)\varphi|\partial_t u|^{r+1}dxds\\
&\le E_\varphi(u)(t)+\tfrac{\lambda}{2}\textstyle\int_t^{t+T}\int_\Omega|\varphi'|\left(|\nabla u(s)|^2+|\partial_t u(s)|^2\right)dxds\\
&+\mu\textstyle\int_t^{t+T}\int_\Omega\left|\varphi'\tfrac{x\cdot\nabla u(s)}{q(x)}\partial_t u(s)\right|dxds+\int_t^{t+T}\int_{S(R)}\varphi\left|\tfrac{x\cdot\nabla u(s)}{R}\partial_t u(s)\right|dSd\tau
\end{aligned}\tag{2.16}$$

Using Young's inequality

$$\int_t^{t+T}\int_{S(R)}\varphi\left|\frac{x\cdot\nabla u}{R}\partial_t u\right|dSd\tau\le\frac{1}{2}\int_t^{t+T}\int_{S(R)}\left(|\partial_r u|^2+|\partial_t u|^2\right)\varphi dSd\tau$$

From (2.15), we infer that

$$\liminf_{R\longrightarrow+\infty}\int_t^{t+T}\int_{S(R)}\left(|\nabla u|^2+|\partial_t u|^2\right)\varphi dSd\tau=0$$

Passing to the limite in (2.16), we get

$$\begin{array}{l}E_\varphi(u)(t+T)+\int_t^{t+T}\int_\Omega a(x)\varphi|\partial_t u|^{r+1}dxds\le E_\varphi(u)(t)\\+\frac{\lambda}{2}\int_t^{t+T}\int_\Omega|\varphi'|\left(|\nabla u(s)|^2+|\partial_t u(s)|^2\right)dxds+\mu\int_t^{t+T}\int_\Omega\left|\varphi'\frac{x\cdot\nabla u}{q(x)}\partial_t u(s)\right|dxds\end{array}$$

Young's inequality, gives

$$\begin{array}{l}E_\varphi(u)(t+T)+\int_t^{t+T}\int_\Omega a(x)(1+\varphi)|\partial_t u|^{r+1}dxds\le E_\varphi(u)(t)\\+\frac{\lambda+\mu}{2}\int_t^{t+T}\int_\Omega|\varphi'|\left(|\nabla u(s)|^2+|\partial_t u(s)|^2\right)dxds.\end{array}$$

□

Next we prove a weighted observability estimate for the local energy of solutions of the system 1.1, the proof is based on the notion of the microlocal defect measures. These measures were introduced by Gérard [14] and Tartar [34] and used to prove stabilization and control results for partial differential equations.

Proposition 2. *We assume that Hyp A holds and* (ω,T) *geometrically controls* Ω*. Let* φ *be a positive function in* $C^2(\mathbb{R}_+)$ *such that* $\varphi'\in L^\infty(\mathbb{R}_+)$*. We suppose that there exists a positive constant* K *such that*

$$\sup_{\mathbb{R}_+}\left|\frac{\varphi''(t)}{\varphi'(t)}\right|\le K.$$

Moreover we assume that the function $t\longmapsto\left|\frac{\varphi'(t)}{\varphi(t)}\right|$ *is decreasing and* $\lim_{t\to+\infty}\left|\frac{\varphi'(t)}{\varphi(t)}\right|=0$*. Let* δ, c_0 *and* c_1 *be a positive constants and* $\chi\in C_0^\infty(\mathbb{R}^d)$*. There exists* $C_{T,\delta,\chi}>0$ *and* $T_1>0$*, such that the following inequality*

$$\begin{array}{l}\int_t^{t+T}\int_\Omega\chi^2(x)\varphi(\mu q(x)+\lambda s)\left(|u|^2+|\nabla u|^2+|\partial_t u|^2\right)dxds\\\le C_{T,\delta,\chi}\int_t^{t+T}\int_\Omega a(x)\varphi(\mu q(x)+\lambda s)|\partial_t u|^2dxds\\+C_{T,\delta,\chi}\int_t^{t+T}\int_\Omega a(x)\varphi(\mu q(x)+\lambda s)|g(t,x)|^2dxds\\+C_{T,\delta,\chi}\int_t^{t+T}\int_\Omega\frac{(\varphi'(\mu q(x)+\lambda s))^2}{\varphi(\mu q(x)+\lambda s)}a(x)|u|^2dxds\\+\delta\int_t^{t+T}\int_\Omega\varphi(\mu q(x)+\lambda s)\left(|\nabla u(s)|^2+|\partial_t u(s)|^2\right)dxds,\end{array}\tag{2.17}$$

holds for every

$$\begin{array}{l}t\ge T_1,\\c_1\le\lambda\le c_0,\\0\le\mu\le c_0,\\g\text{ such that }\sqrt{\varphi}g\in L^2_{loc}\left(\mathbb{R}_+,L^2(\Omega)\right),\end{array}$$

for all

$$u\in C^0\left(\mathbb{R}_+,H_0^1(\Omega)\right)\cap C^1\left(\mathbb{R}_+,L^2(\Omega)\right)$$

solution of

$$\left\{\begin{array}{ll} \partial_t^2 u-\Delta u=g & \textit{in } \mathbb{R}_+\times\Omega, \\ u=0 & \textit{on } \mathbb{R}_+\times\Gamma, \\ u(0,x)=u_0 \quad \textit{and} \quad \partial_t u(0,x)=u_1. & \end{array}\right. \tag{2.18}$$

such that $E_\varphi(u)(0)<\infty$.

Proof. To prove this result we argue by contradiction: If (2.17) was false, there would exist a sequences

$$\begin{array}{l} (t_n) \text{ such that } t_n \underset{n\to+\infty}{\longrightarrow} +\infty \\ (\lambda_n) \text{ such that } c_1\le\lambda_n\le c_0 \\ (\mu_n) \text{ such that } 0\le\mu_n\le c_0 \\ (g_n) \text{ such that } \sqrt{\varphi}g_n\in L^2_{loc}\left(\mathbb{R}_+,L^2(\Omega)\right) \end{array}$$

and a sequence of solutions (u_n) in $C^0\left(\mathbb{R}_+,H_0^1(\Omega)\right)\cap C^1\left(\mathbb{R}_+,L^2(\Omega)\right)$ with $E_\varphi(u_n)(0)<\infty$ and such that

$$\begin{array}{l} \int_{t_n}^{t_n+T}\int_\Omega \chi^2(x)\varphi(\mu_n q(x)+\lambda_n s)\left(|u_n|^2+|\nabla u_n|^2+|\partial_t u_n|^2\right)dxds \\ \ge n\left(\int_{t_n}^{t_n+T}\int_\Omega a(x)\varphi(\mu_n q(x)+\lambda_n s)|\partial_t u_n|^2 dxds\right) \\ +n\int_{t_n}^{t_n+T}\int_\Omega \varphi(\mu_n q(x)+\lambda_n s)|g_n(s,x)|^2 dxds \\ +n\left(\int_{t_n}^{t_n+T}\int_\Omega a(x)\frac{(\varphi'(\mu_n q(x)+\lambda_n s))^2}{\varphi(\mu_n q(x)+\lambda_n s)}|u_n|^2 dxds\right) \\ +\delta\int_{t_n}^{t_n+T}\int_\Omega \varphi(\mu_n q(x)+\lambda_n s)\left(|\nabla u(t)|^2+|\partial_t u(t)|^2\right)dxds \end{array} \tag{2.19}$$

Setting

$$\begin{array}{l} \sigma_n^2=\int_{t_n}^{t_n+T}\int_\Omega \chi^2(x)\varphi(\mu_n q(x)+\lambda_n s)\left(|u_n|^2+|\nabla u_n|^2+|\partial_t u_n|^2\right)dxds \\ \text{and } v_n=\frac{(\varphi(\mu_n q(x)+\lambda_n s))^{\frac{1}{2}}u_n(t_n+\cdot)}{\sigma_n}. \end{array}$$

Therefore from (2.19) we infer that

$$\frac{1}{\sigma_n^2}\int_{t_n}^{t_n+T}\int_\Omega \varphi(\mu_n q(x)+\lambda_n s)\left(|\nabla u(t)|^2+|\partial_t u(t)|^2\right)dxdt\le\frac{1}{\delta}, \tag{2.20}$$

and

$$\begin{array}{l} \frac{1}{\sigma_n^2}\int_{t_n}^{t_n+T}\int_\Omega a(x)\varphi(\mu_n q(x)+\lambda_n s)|\partial_t u_n|^2 dxds \underset{n\to+\infty}{\longrightarrow} 0 \\ \frac{1}{\sigma_n^2}\int_{t_n}^{t_n+T}\int_\Omega \varphi(\mu_n q(x)+\lambda_n s)|g_n(s,x)|^2 dxds \underset{n\to+\infty}{\longrightarrow} 0 \\ \frac{1}{\sigma_n^2}\int_{t_n}^{t_n+T}\int_\Omega a(x)\frac{(\varphi'(\mu_n q(x)+\lambda_n s))^2}{\varphi(\mu_n q(x)+\lambda_n s)}|u_n|^2 dxds \underset{n\to+\infty}{\longrightarrow} 0 \end{array}. \tag{2.21}$$

It is clear that v_n is a solution of the following system

$$\left\{\begin{array}{ll} \partial_t^2 v_n-\Delta v_n=f_n(t,x) & \text{in } \mathbb{R}_+\times\Omega, \\ v_n(t,x)=0 & \text{on } \mathbb{R}_+\times\Gamma, \\ (v_{n,0},v_{n,1})\in H_0^1(\Omega)\times L^2(\Omega), & \end{array}\right.$$

with

$$\begin{aligned}
&f_n(t,x)=\tfrac{\mu_n}{2\sigma_n}\left[\mu_n\left(\varphi''(\varphi)^{-\frac{1}{2}}-\tfrac{1}{2}(\varphi')^2\varphi^{-3/2}\right)\tfrac{|x|^2}{q^2}+\left(\tfrac{d}{q}-\tfrac{|x|^2}{q^3}\right)\varphi'(\varphi)^{-\frac{1}{2}}\right]u_n(t_n+t)\\
&+\tfrac{\lambda_n}{2\sigma_n}\left[\lambda_n\left(\varphi''(\varphi)^{-\frac{1}{2}}-\tfrac{1}{2}(\varphi')^2\varphi^{-3/2}\right)\right]u_n(t_n+t)\\
&+\tfrac{\varphi'(\varphi)^{-\frac{1}{2}}}{\sigma_n}\left(\lambda_n\partial_t u_n(t_n+t)+\mu_n\tfrac{x\cdot\nabla u_n(t_n+t)}{q}\right)-\tfrac{1}{\sigma_n}\varphi^{\frac{1}{2}}g_n(t_n+t,x)
\end{aligned}$$

It is easy to see that (2.19), gives

$$\begin{aligned}
&\textstyle\int_0^T\int_\Omega\left|\tfrac{\mu_n}{2\sigma_n}\left[\mu_n\left(\varphi''(\varphi)^{-\frac{1}{2}}-\tfrac{1}{2}(\varphi')^2\varphi^{-3/2}\right)\tfrac{|x|^2}{q^2}+\left(\tfrac{d}{q}-\tfrac{|x|^2}{q^3}\right)\varphi'(\varphi)^{-\frac{1}{2}}\right]u_n(t_n+t)\right|^2dxdt\\
&\le\tfrac{C}{\sigma_n^2}\textstyle\int_{t_n}^{t_n+T}\int_\Omega a(x)\tfrac{(\varphi'(\mu q(x)+\lambda s))^2}{\varphi(\mu q(x)+\lambda s)}|u_n|^2dxds\underset{n\to+\infty}{\longrightarrow}0
\end{aligned}$$

and

$$\begin{aligned}
&\textstyle\int_0^T\int_\Omega\left|\tfrac{\lambda_n}{2\sigma_n}\left[\lambda\left(\varphi''(\varphi)^{-\frac{1}{2}}-\tfrac{1}{2}(\varphi')^2\varphi^{-3/2}\right)\right]u_n(t_n+t)\right|^2dxdt\\
&\le\tfrac{C}{\sigma_n^2}\textstyle\int_{t_n}^{t_n+T}\int_\Omega a(x)\tfrac{(\varphi'(\mu q(x)+\lambda s))^2}{\varphi(\mu q(x)+\lambda s)}|u_n|^2dxds\underset{n\to+\infty}{\longrightarrow}0.
\end{aligned}$$

Moreover we have,

$$\begin{aligned}
&\textstyle\int_0^T\int_\Omega\left|\tfrac{\varphi'(\varphi)^{-\frac{1}{2}}}{\sigma_n}\left(\lambda_n\partial_t u_n(t_n+t)+\mu\tfrac{x\cdot\nabla u(t_n+t)}{q}\right)\right|^2dxdt\\
&\le\tfrac{C}{\sigma_n^2}\left(\tfrac{\varphi'(c_1t_n)}{\varphi(c_1t_n)}\right)^2\textstyle\int_0^T\int_\Omega\varphi(\mu_nq(x)+\lambda_n(t_n+s))\left(|\partial_tu_n(t_n+t)|^2+|\nabla u_n(t_n+t)|^2\right)dxdt\\
&\le\tfrac{C}{\delta\sigma_n^2}\left(\tfrac{\varphi'(c_1t_n)}{\varphi(c_1t_n)}\right)^2\underset{n\to+\infty}{\longrightarrow}0,
\end{aligned}$$

and

$$\frac{1}{\sigma_n^2}\int_0^T\int_\Omega a(x)\varphi(\mu_nq(x)+\lambda_n(t_n+s))|g_n(t_n+t,x)|^2dxds\underset{n\to+\infty}{\longrightarrow}0.$$

We conclude that

$$\int_0^T\int_\Omega|f_n(s,x)|^2dxds\underset{n\to+\infty}{\longrightarrow}0.\tag{2.22}$$

On the other hand, we have

$$\begin{aligned}
&\textstyle\int_0^T E_{v_n}(t)\,dt\le\tfrac{1}{\sigma_n^2}\int_{t_n}^{t_n+T}\int_\Omega\varphi(\mu_nq(x)+\lambda_ns)\left(|\nabla u_n(t)|^2+|\partial_tu_n(t)|^2\right)dxdt\\
&+\tfrac{\lambda_n^2+\mu_n^2}{\sigma_n^2}\textstyle\int_{t_n}^{t_n+T}\int_\Omega\tfrac{(\varphi'(\mu_nq(x)+\lambda_ns))^2}{\varphi(\mu_nq(x)+\lambda_ns)}|u_n(t)|^2dxdt\\
&\le\tfrac{1}{\sigma_n^2}\textstyle\int_{t_n}^{t_n+T}\int_\Omega\varphi(\mu_nq(x)+\lambda_ns)\left(|\nabla u_n(t)|^2+|\partial_tu_n(t)|^2\right)dxdt\\
&+\tfrac{\lambda_n^2+\mu_n^2}{\sigma_n^2\epsilon_0}\textstyle\int_{t_n}^{t_n+T}\int_\Omega a(x)\tfrac{(\varphi'(\mu_nq(x)+\lambda_ns))^2}{\varphi(\mu_nq(x)+\lambda_ns)}|u_n(t)|^2dxdt\\
&+2C_Lc_0^2\left(\tfrac{\varphi'(c_1t_n)}{\varphi(c_1t_n)}\right)^2\textstyle\int_0^T\int_\Omega|\nabla v_n(t)|^2dxdt
\end{aligned}$$

Using the fact that

$$\lim_{n\to+\infty}\left|\frac{\varphi'(t_n)}{\varphi(t_n)}\right|=0$$

we infer that, there exists $n_0 \in \mathbb{N}$, such that

$$\int_0^T E_{v_n}(t)\,dt \leq C_\delta \text{ for all } n \geq n_0$$

Now we prove that the total energy of v_n is bounded on $[0,T]$. Let Z_n be the solution of the following system

$$\begin{cases} \partial_t^2 Z_n - \Delta Z_n = 0 & \mathbb{R}_+ \times \Omega, \\ Z_n = 0 & \mathbb{R}_+ \times \Gamma, \\ (Z_n(0), \partial_t Z_n(0)) = (v_n(0), \partial_t v_n(0)). \end{cases} \tag{2.23}$$

The hyperbolic energy inequality gives

$$\sup_{[0,T]} E_{v_n - Z_n}(s) \leq C_T \left\| f_n(t,x) \right\|^2_{L^2([0,T],L^2(\Omega))}$$

Now using (2.22), we deduce that

$$\sup_{[0,T]} E_{v_n - Z_n}(s) \underset{n\to+\infty}{\longrightarrow} 0.$$

Turn into account of the estimate above, we obtain

$$\begin{aligned} TE_{v_n}(0) &= TE_{Z_n}(0) = \int_0^T E_{Z_n}(t)\,dt \\ &\leq 2T\sup_{[0,T]} E_{v_n - Z_n}(s) + 2\int_0^T E_{v_n}(t)\,dt \\ &\leq C_{T,\delta} \end{aligned}$$

which means

$$\sup_{[0,T]} E_{Z_n}(s) = E_{v_n}(0) \leq C_{T,\delta}.$$

Using the result above and ([11, proposition 2]), we infer that

$$\begin{aligned} &E_{v_n}(t) \leq 2e^T \left(E_{v_n}(0) + \textstyle\int_0^T \int_\Omega |f_n(\tau,x)|^2\,dxd\tau \right) \text{ for } 0 \leq t \leq T \\ &\leq C_{T,\delta}, \text{ for all } 0 \leq t \leq T. \end{aligned}$$

We have

$$\begin{aligned} &\textstyle\int_0^T \int_\Omega a(x) |\partial_t v_n|^2\,dxdt \leq \frac{\lambda_n^2}{\sigma_n^2} \int_{t_n}^{t_n+T} \int_\Omega a(x) \frac{(\varphi'(\mu_n q(x)+\lambda_n s))^2}{\varphi(\mu_n q(x)+\lambda_n s)} |u_n(s)|^2\,dxds \\ &+\textstyle\frac{2}{\sigma_n^2} \int_{t_n}^{t_n+T} \int_\Omega a(x)\varphi(\mu_n q(x) + \lambda_n s) |\partial_t u_n|^2\,dxds. \end{aligned}$$

Now using (2.21), we deduce that

$$\int_0^T \int_\Omega a(x) |\partial_t v_n|^2\,dxdt \underset{n\to+\infty}{\longrightarrow} 0. \tag{2.24}$$

Using the microlocal defect measures associated to solutions (2.23) and proceeding as in [12, Proof of proposition 3] by taking into account of (2.22) and (2.24), we infer that

$$\int_0^T \int_\Omega \chi^2(x) \left(|v_n|^2 + |\nabla v_n|^2 + |\partial_t v_n|^2 \right) dxds \underset{n\to+\infty}{\longrightarrow} 0.$$

The contradiction follows from the fact that

$$
\begin{aligned}
&1 \leq C \int_0^T \int_\Omega \chi^2(x) \left(|v_n|^2 + |\nabla v_n|^2 + |\partial_t v_n|^2 \right) dxds \\
&+ \tfrac{C}{\sigma_n^2} \int_{t_n}^{t_n+T} \int_\Omega \tfrac{(\varphi'(\mu_n q(x)+\lambda_n s))^2}{\varphi(\mu_n q(x)+\lambda_n s)} a(x) |u_n|^2 \, dxds + C \left(\tfrac{\varphi'(c_1 t_n)}{\varphi(c_1 t_n)} \right)^2 \int_0^T \int_\Omega |\nabla v_n(t)|^2 \, dxdt \underset{n\to+\infty}{\longrightarrow} 0.
\end{aligned}
$$

□

3. Proof of Theorem 1

3.1. **Preliminary results.**

Proposition 3. *We assume that Hyp A holds and (ω,T) geometrically controls Ω. Let $\beta > -1$. Setting*

$$f(s) = \frac{\ln^\beta(b+s)}{b+s}, \ \ f_1(s) = \frac{\ln^\beta(b+s)}{(b+s)^2}$$

where

$$\ln b = \max \left((2(r+1))^{r+1}, \beta, \tfrac{\beta + \sqrt{|\beta^2+4\beta|}}{2}, \tfrac{\beta+1-r}{r-1}, \left(\tfrac{8(r+1)(\beta+1)}{1-\delta_0} \right)^{r+1} \right)$$

Let δ,$R_0 > 0$. There exists $C_{T,\delta,R_0} > 0$ and $T_1 > 0$, such that the following inequality

$$
\begin{aligned}
&\int_t^{t+T} \int_{\Omega\cap B_{R_0}} f(q(x)+s) \left(|\nabla u|^2 + |\partial_t u|^2 \right) dxds \\
&\leq C_{T,\delta,R_0} \int_t^{t+T} \int_\Omega a(x) f(q(x)+s) \left(|\partial_t u|^2 + |\partial_t u|^{2r} \right) dxds \\
&+ C_{T,\delta,R_0} \int_t^{t+T} \int_\Omega a(x) f_1'(q(x)+s) |u|^2 \, dxds \\
&+ \delta \int_t^{t+T} \int_\Omega f(q(x)+s) \left(|\nabla u(s)|^2 + |\partial_t u(s)|^2 \right) dxds,
\end{aligned}
\tag{3.1}
$$

holds for every $t \geq T_1$ and for all u solution of (1.1) with initial data (u_0,u_1) in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$.

Proof. It is clear that there exists a positive constant K, such that

$$\sup_{\mathbb{R}_+} \left| \frac{f''(t)}{f'(t)} \right| \leq K.$$

In addition the function $t \longmapsto \left| \frac{f'(t)}{f(t)} \right|$ is decreasing and $\lim\limits_{t\to+\infty} \left| \frac{f'(t)}{f(t)} \right| = 0$. Moreover there exists $C > 0$, such that

$$\frac{(f'(t))^2}{f(t)} \leq C \ \left(-f_1'(t) \right), \text{ for all } t \geq 0.$$

Since

$$\partial_t u \in L^\infty \left(\mathbb{R}_+, H_0^1(\Omega) \right).$$

From Sobolev imbedding, we deduce that

$$\sqrt{af} \, |\partial_t u|^r \in L^2_{loc} \left(\mathbb{R}_+, L^2(\Omega) \right).$$

Let $\chi \in C_0^\infty \left(\mathbb{R}^d \right)$ such that

$$\chi(x) = \begin{cases} 1 & \text{for } |x| \leq R_0 \\ 0 & \text{for } |x| \geq 2R_0 \end{cases}.$$

Therefore using (2.17), we infer that

$$\begin{aligned}
&\int_t^{t+T}\int_{\Omega\cap B_{R_0}} f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&\leq \int_t^{t+T}\int_\Omega \chi^2\left(x\right) f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&\leq C_{T,\delta,R_0}\int_t^{t+T}\int_\Omega a\left(x\right) f\left(q\left(x\right)+s\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\\
&-CC_{T,\delta,R_0}\int_t^{t+T}\int_\Omega a\left(x\right) f_1'\left(q\left(x\right)+s\right)\left|u\right|^2 dxds\\
&+\delta\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\left(s\right)\right|^2+\left|\partial_t u\left(s\right)\right|^2\right)dxds,
\end{aligned}$$

This finishes the proof of the proposition. □

In order to prove theorem 1 we need the following result.

Lemma 1. *Let* $\beta>-1$, $T>0$ *and* $0<\delta_0<1$. *Setting*

$$\varphi\left(s\right)=\ln^{\beta+1}\left(b+s\right),\ f\left(s\right)=\tfrac{\ln^{\beta}(b+s)}{b+s},\ f_1\left(s\right)=\tfrac{\ln^{\beta}(b+s)}{(b+s)^2}$$
$$\text{and } f_2\left(s\right)=\tfrac{\ln^{\beta-r+1}(b+s)}{(b+s)^r},$$

with

$$\ln b=\max\left(\left(2\left(r+1\right)\right)^{r+1},\beta,\tfrac{\beta+\sqrt{\left|\beta^2+4\beta\right|}}{2},\tfrac{\beta+1-r}{r-1},\left(\tfrac{8(r+1)(\beta+1)}{1-\delta_0}\right)^{r+1}\right)$$

Setting $w=\psi u$ *and* $v=\left(1-\psi\right)u$ *where* u *is a solution of* (1.1) *with initial data in* $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$ *such that* $E_\varphi\left(u\right)\left(0\right)<\infty$. *Let* $t\geq 0$ *and we set*

$$\begin{aligned}
&X\left(t\right)=\int_\Omega f\left(q\left(x\right)+t\right)v\left(t\right)\partial_t v\left(t\right)dx+\tfrac{k_1}{2}\int_\Omega f_1\left(q\left(x\right)+t\right)a\left(x\right)\left|u\left(t\right)\right|^2dx\\
&+\int_\Omega a\left(x\right)f_2\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^{r+1}dx+\tfrac{k}{2}\int_\Omega \ln^{\beta+1}\left(b+q\left(x\right)+t\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dx,
\end{aligned}$$

where

$$k=\frac{1-\delta_0}{2\left(\beta+1\right)},\ k_1>0$$

We have

$$\begin{aligned}
&X\left(t+T\right)-X\left(t\right)+\tfrac{\delta_0}{2}\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&-\left(\tfrac{k_1}{4}-\tfrac{2}{\epsilon_0}\right)\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\left(t\right)\right|^2dxds\\
&-\tfrac{1}{2}\int_t^{t+T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\left(t\right)\right|^{r+1}dxds\\
&+\tfrac{1-\delta_0}{4(\beta+1)}\int_t^{t+T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\\
&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}} f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4(1+|\beta|)}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left|\partial_t u\right|^2dxds
\end{aligned}\tag{3.2}$$

for some $C>0$.

Proof. We have

$$\int_\Omega \varphi\left(q\left(x\right)\right)\left(\left|\nabla u_0\right|^2+\left|u_1\right|^2\right)dx<\infty.$$

Then from (2.2), we infer

$$\begin{aligned}&E_{\varphi}(u)(t+T)+\int_{t}^{t+T}\int_{\Omega}a(x)\varphi(q(x)+s)\left|\partial_{t}u\right|^{r+1}dxds\\&\leq E_{\varphi}(u)(t)+(\beta+1)\int_{t}^{t+T}\int_{\Omega}f(q(x)+s)\left(\left|\nabla u(s)\right|^{2}+\left|\partial_{t}u(s)\right|^{2}\right)dxds.\end{aligned}\tag{3.3}$$

Setting

$$\begin{aligned}&X_{0}(t)=\int_{\Omega}f(q(x)+t)v(t)\partial_{t}v(t)dx+\frac{k_{1}}{2}\int_{\Omega}f_{1}(q(x)+t)a(x)\left|u(t)\right|^{2}dx\\&+\int_{\Omega}a(x)f_{2}(q(x)+t)\left|u(t)\right|^{r+1}dx\end{aligned}$$

We have $v=(1-\psi)u$. Then v is a solution of

$$\begin{cases}\partial_{t}^{2}v-\Delta v+a(x)\left|\partial_{t}u\right|^{r-1}\partial_{t}v=g(t,x) & \mathbb{R}_{+}\times\Omega,\\ v=0 & \mathbb{R}_{+}\times\Gamma,\\ (v(0),\partial_{t}v(0))=(1-\psi)(u_{0},u_{1}),\end{cases}\tag{3.4}$$

with

$$g(t,x)=2\nabla\psi\nabla u+u\Delta\psi.$$

Now using the fact that v is a solution of (3.4), we get

$$\begin{aligned}&\frac{d}{dt}X_{0}(t)=\int_{\Omega}\left(\left|\partial_{t}v(t)\right|^{2}-\left|\nabla v(t)\right|^{2}-a(x)\left|\partial_{t}u(t)\right|^{r-1}v\partial_{t}v(t)+g(t,x)v\right)f(q(x)+t)dx\\&+\int_{\Omega}f'(q(x)+t)v(t)\partial_{t}v(t)dx+\frac{1}{2}\int_{\Omega}\Delta f(q(x)+t)\left|v(t)\right|^{2}dx\\&k_{1}\left(\int_{\Omega}a(x)f_{1}(q(x)+t)u(t)\partial_{t}u(t)dx+\frac{1}{2}\int_{\Omega}a(x)f_{1}'(q(x)+t)\left|u(t)\right|^{2}dx\right)\\&+\int_{\Omega}a(x)f_{2}'(q(x)+t)\left|u\right|^{r+1}dx+(r+1)\int_{\Omega}a(x)f_{2}(q(x)+t)\left|u\right|^{r-1}u\partial_{t}udx\end{aligned}\tag{3.5}$$

We have

$$\begin{aligned}&\frac{(f'(s))^{2}}{f(s)}\leq(1+|\beta|)\frac{\ln^{\beta}(b+s)}{(b+s)^{3}}\leq-(1+|\beta|)f_{1}'(s)\\&\text{and}\\&\frac{(f_{1}(s))^{2}}{f(s)}=\frac{\ln^{\beta}(b+s)}{(b+s)^{3}}\leq-f_{1}'(s)\end{aligned}$$

By taking into account that the support of $(1-\psi)$ is contained in the set $\{x\in\Omega,a(x)>\epsilon_{0}\}$ and using Young's inequality we infer that

$$\begin{aligned}&\left|\int_{\Omega}f'(q(x)+t)v(t)\partial_{t}v(t)dx\right|\\&\leq-\frac{k_{1}}{8}\int_{\Omega}a(x)f_{1}'(q(x)+t)\left|u(t)\right|^{2}dx+\frac{8(1+|\beta|)}{\epsilon_{0}^{2}k_{1}}\int_{\Omega}a(x)f(q(x)+t)\left|\partial_{t}u(t)\right|^{2}dx\end{aligned}$$

and

$$\begin{aligned}&\left|k_{1}\int_{\Omega}af_{1}(q(x)+t)u(t)\partial_{t}u(t)dx\right|\\&\leq-\frac{k_{1}}{8}\int_{\Omega}a(x)f_{1}'(q(x)+t)\left|u(t)\right|^{2}dx+8k_{1}\int_{\Omega}a(x)f(q(x)+t)\left|\partial_{t}u(t)\right|^{2}dx\end{aligned}$$

Using Young's inequality we deduce that

$$k(\beta+1)\int_{\Omega}f(q(x)+t)\frac{x\cdot\nabla u}{q(x)}\partial_{t}u(t)dx\leq\frac{k(\beta+1)}{2}\int_{\Omega}f(q(x)+t)\left(\left|\nabla u\right|^{2}+\left|\partial_{t}u\right|^{2}\right)dx$$

We have

$$f'(s) = \beta \tfrac{\ln^{\beta-1}(b+s)}{(b+s)^2} - \tfrac{\ln^{\beta}(b+s)}{(b+s)^2} \leq 0$$

and

$$\begin{aligned} & f''(s) = 2\tfrac{\ln^{\beta}(b+s)}{(b+s)^3} - 3\beta \tfrac{\ln^{\beta-1}(b+s)}{(b+s)^3} + \beta \left(\ln^{\beta-2}(b+s)\right) \tfrac{\beta-1}{(b+s)^3} \\ & \leq 4\tfrac{\ln^{\beta}(b+s)}{(b+s)^3} \leq -4f_1'(s) \end{aligned}$$

Since

$$\begin{aligned} & \Delta f(q(x)+t) = f''((q(x)+t)) \tfrac{|x|^2}{q^2} + f'((q(x)+t)) \left(\tfrac{d}{q} - \tfrac{|x|^2}{q^3}\right) \\ & \leq f''((q(x)+t)) \end{aligned}$$

Therefore we obtain

$$\tfrac{1}{2} \int_\Omega \Delta f(q(x)+t) |v(t)|^2 \, dx \leq -\tfrac{2}{\epsilon_0} \int_\Omega a(x) f_1'(q(x)+t) |u(t)|^2 \, dx$$

We have

$$\begin{aligned} & -f_2'(s) \geq \tfrac{\ln^{\beta-r+1}(b+s)}{(b+s)^{r+1}} \\ & (f(s))^{r+1} \ln^{-r(\beta+1)}(s) \leq \tfrac{-f_2'(s)}{\ln(b+s)} \\ & (f_2(s))^{\frac{r+1}{r}} \ln^{-\frac{\beta+1}{r}}(s) \leq \tfrac{-f_2'(s)}{\ln(b+s)} \end{aligned}$$

Using Holder's and Young's inequalities, we obtain

$$\begin{aligned} & \int_\Omega a(x) f(q(x)+s) |\partial_t u(t)|^{r-1} u \partial_t u dx \\ & \leq (\ln b)^{-\frac{1}{r+1}} \left(\int_\Omega a(x) \ln^{\beta+1}(q(x)+s) |\partial_t u|^{r+1} dx\right)^{\frac{r}{r+1}} \times \left(-\int_\Omega a(x) f_2'(q(x)+s) |u|^{r+1} dx\right)^{\frac{1}{r+1}} \\ & \leq (\ln b)^{-\frac{1}{r+1}} \int_\Omega a(x) \ln^{\beta+1}(q(x)+s) |\partial_t u|^{r+1} dx - (\ln b)^{-\frac{1}{r+1}} \int_\Omega a(x) f_2'(q(x)+s) |u|^{r+1} dx \end{aligned}$$

and

$$\begin{aligned} & (r+1) \int_\Omega a(x) f_2(q(x)+s) |u|^{r-1} u \partial_t u dx \\ & \leq (r+1) (\ln b)^{-\frac{r}{r+1}} \left(\int_\Omega a(x) \ln^{\beta+1}(b+q(x)+s) |\partial_t u|^{r+1} dx\right)^{\frac{1}{r+1}} \\ & \times \left(-\int_\Omega a(x) f_2'(q(x)+s) |u|^{r+1} dx\right)^{\frac{r}{r+1}} \\ & \leq (\ln b)^{-\frac{1}{r+1}} \int_\Omega a(x) \ln^{\beta+1}(q(x)+s) |\partial_t u|^{r+1} dx - r(\ln b)^{-\frac{1}{r+1}} \int_\Omega a(x) f_2'(q(x)+s) |u|^{r+1} dx \end{aligned}$$

Thus

$$\begin{aligned} & \int_t^{t+T} \int_\Omega a(x) f(q(x)+s) |\partial_t u|^{r-1} u \partial_t u dx ds + (r+1) \int_t^{t+T} \int_\Omega a(x) f_2(q(x)+s) |u|^{r-1} u \partial_t u dx ds \\ & \leq (r+1) (\ln b)^{-\frac{1}{r+1}} \int_t^{t+T} \int_\Omega a(x) \ln^{\beta+1}(b+q(x)+s) |\partial_t u|^{r+1} dx ds \\ & -(r+1) (\ln b)^{-\frac{1}{r+1}} \int_t^{t+T} \int_\Omega a(x) f_2'(q(x)+s) |u|^{r+1} dx ds \end{aligned}$$

Since $a(x) > \epsilon_0$ on the support of $(1-\psi)$, therefore

$$\begin{aligned}
&\textstyle\int_\Omega f(q(x)+t)\left(|\partial_t v(t)|^2 - |\nabla v(t)|^2\right)dx \\
&= 2\textstyle\int_\Omega f(q(x)+t)\,|\partial_t v(t)|^2\,dx - \int_\Omega f(q(x)+t)\left(|\nabla v|^2 + |\partial_t v|^2\right)dx \\
&\leq 2\textstyle\int_\Omega f(q(x)+t)\,|\partial_t v(t)|^2\,dx - \frac{1}{2}\int_\Omega f(q(x)+t)\left(|\nabla u|^2 + |\partial_t u|^2\right)dx \\
&+ \textstyle\int_\Omega f(q(x)+t)\left(|\nabla w|^2 + |\partial_t w|^2\right)dx \\
&\leq \tfrac{2}{\epsilon_0}\textstyle\int_\Omega a(x) f(q(x)+t)\,|\partial_t u(t)|^2\,dx - \frac{1}{2}\int_\Omega f(q(x)+t)\left(|\nabla u|^2 + |\partial_t u|^2\right)dx \\
&+ \textstyle\int_\Omega f(q(x)+t)\left(|\nabla w|^2 + |\partial_t w|^2\right)dx
\end{aligned}$$

The function $\chi \in C_0^\infty(\mathbb{R}^d)$ and support of ψ is contained in B_{2L}. Then it is clear that there exists a positive constant C, such that

$$\left|\textstyle\int_\Omega f(q(x)+t)\,g(t,x)\,v\,dx\right| \leq C\textstyle\int_{\Omega\cap B_{2L}} f(q(x)+t)\left(|u|^2 + |\nabla u|^2\right)dx.$$

Since $w = \psi u$ Therefore, there exists a positive constant C such that

$$\begin{aligned}
&\textstyle\int_\Omega f(q(x)+s)\left(|\nabla w(s)|^2 + |\partial_t w(s)|^2\right)dxds \\
&\leq C\textstyle\int_{\Omega\cap B_{2L}} f(q(x)+s)\left(|u|^2 + |\nabla u|^2 + |\partial_t u|^2\right)dxds
\end{aligned}$$

By taking into account of the estimates above and integrating (3.5) between t and $t+T$, we obtain

$$\begin{aligned}
&X(t+T) - X(t) + \tfrac{(1-2(1+\beta)k)}{2}\textstyle\int_t^{t+T}\int_\Omega f(q(x)+s)\left(|\nabla u|^2 + |\partial_t u|^2\right)dxds \\
&-\left(\tfrac{k_1}{4} - \tfrac{2}{\epsilon_0}\right)\textstyle\int_t^{t+T}\int_\Omega a(x) f_1'(q(x)+s)\,|u(s)|^2\,dxds \\
&-\left(1-(r+1)(\ln b)^{-\frac{1}{r+1}}\right)\textstyle\int_t^{t+T}\int_\Omega a(x) f_2'(q(x)+s)\,|u|^{r+1}\,dxds \\
&+\left(k-(r+1)(\ln b)^{-\frac{1}{r+1}}\right)\textstyle\int_t^{t+T}\int_\Omega a(x)\ln^{\beta+1}(b+q(x)+s)\,|\partial_t u|^{r+1}\,dxds \\
&\leq C\textstyle\int_t^{t+T}\int_{\Omega\cap B_{2L}} f(q(x)+s)\left(|u|^2 + |\nabla u|^2\right)dxds \\
&+\left(\tfrac{2}{\epsilon_0} + \tfrac{8(1+|\beta|)}{\epsilon_0^2 k_1} + 8k_1\right)\textstyle\int_t^{t+T}\int_\Omega a(x) f(q(x)+s)\,|\partial_t u|^2\,dxds.
\end{aligned}$$

Using the fact that $k = \frac{1-\delta_0}{2(\beta+1)}$ and

$$\ln b \geq \max\left((2(r+1))^{r+1}, \left(\frac{4(r+1)(\beta+1)}{1-\delta_0}\right)^{r+1}\right)$$

We obtain

$$\begin{aligned}
&X\left(t+T\right)-X\left(t\right)+\tfrac{\delta_0}{2}\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&-\left(\tfrac{k_1}{4}-\tfrac{2}{\epsilon_0}\right)\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\left(t\right)\right|^2dxds\\
&-\tfrac{1}{2}\int_t^{t+T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds\\
&+\tfrac{1-\delta_0}{4(\beta+1)}\int_t^{t+T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\\
&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}}f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4(1+|\beta|)}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+t\right)\left|\partial_t u\right|^2dxds.
\end{aligned}\tag{3.6}$$

□

3.2. **Proof of Theorem 1.** We assume that Hyp A holds and (ω,T_0) geometrically controls Ω. Let u be a solution of (1.1) with initial data in $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$ such that $E_\varphi\left(u\right)\left(0\right)<\infty$. Let $0<\delta_0\ll 1$ and $T\geq\max\left(T_0,T_1\right)$. Setting

$$\varphi\left(s\right)=\ln^{\beta+1}\left(b+s\right),\ f\left(s\right)=\tfrac{\ln^\beta(b+s)}{b+s},\ f_1\left(s\right)=\tfrac{\ln^\beta(b+s)}{(b+s)^2}\text{ and }f_2\left(s\right)=\tfrac{\ln^{\beta-r+1}(b+s)}{(b+s)^r}$$

with

$$\ln b=\max\left(\left(2\left(r+1\right)\right)^{r+1},\beta,\tfrac{\beta+\sqrt{\left|\beta^2+4\beta\right|}}{2},\tfrac{\beta+1-r}{r-1},\left(\tfrac{4(r+1)(\beta+1)}{1-\delta_0}\right)^{r+1}\right)$$

and

$$\begin{array}{ll}\gamma=\beta+1>0 & \text{if } 1<r<1+\frac{2}{d}\\ 0<\gamma=\beta+1<\frac{2}{r-1} & \text{if } r=1+\frac{2}{d}\end{array}$$

We define

$$\begin{aligned}
&X\left(t\right)=\int_\Omega f\left(q\left(x\right)+t\right)v\left(t\right)\partial_t v\left(t\right)dx+\tfrac{k_1}{2}\int_\Omega f_1\left(q\left(x\right)+t\right)a\left(x\right)\left|u\left(t\right)\right|^2dx\\
&+\int_\Omega a\left(x\right)f_2\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^{r+1}dx+\tfrac{k}{2}\int_\Omega\ln^{\beta+1}\left(b+q\left(x\right)+t\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dx
\end{aligned}$$

where $v=\left(1-\psi\right)u$ and

$$k=\frac{1-\delta_0}{2\left(\beta+1\right)},\ k_1>0.$$

According to lemma 1,

$$\begin{aligned}
&X\left(t+T\right)-X\left(t\right)+\tfrac{\delta_0}{2}\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&-\left(\tfrac{k_1}{4}-\tfrac{2}{\epsilon_0}\right)\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds\\
&-\tfrac{1}{2}\int_t^{t+T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\left(t\right)\right|^{r+1}dxds\\
&+\tfrac{1-\delta_0}{4(\beta+1)}\int_t^{t+T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\\
&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}}f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4(1+|\beta|)}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left|\partial_t u\right|^2dxds
\end{aligned}\tag{3.7}$$

The estimate (4.1), gives

$$\begin{aligned}&C\int_t^{t+T}\int_{\Omega\cap B_{2L}} f\left(q\left(x\right)+s\right)\left(\left|u\right|^2+\left|\nabla u\right|^2\right)dxds\\&\leq C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\\&-C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds\\&+C\delta\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds,\ \text{for all } t\geq T_1\end{aligned}$$

(3.7) and the estimates above gives

$$\begin{aligned}&X\left(t+T\right)-X\left(t\right)+\left(\tfrac{\delta_0}{2}-C\delta\right)\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&-\left(\tfrac{k_1}{4}-\tfrac{2}{\epsilon_0}-C_{T,\delta}\right)\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds\\&-\tfrac{1}{2}\int_t^{t+T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds\\&+\tfrac{1-\delta_0}{4(\beta+1)}\int_t^{t+T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\\&\leq 2\left(\tfrac{1}{\epsilon_0}+\tfrac{4(1+|\beta|)}{\epsilon_0^2k_1}+4k_1+C_{T,\delta}\right)\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left|\partial_t u\right|^2dxds\\&+C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left|\partial_t u\right|^{2r}dxds\end{aligned}\tag{3.8}$$

for every $t\geq T$. Using Young's inequality we get

$$\begin{aligned}X\left(t\right)&\leq\left(\tfrac{k_1}{2}+\tfrac{1}{2\epsilon_0\epsilon}\right)\int_\Omega a\left(x\right)f_1\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^2dx\\&+\left(k+\tfrac{\epsilon}{\epsilon_0}\right)\int_\Omega\ln^{\beta+1}\left(b+q\left(x\right)+t\right)\left(\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx\\&+\int_\Omega a\left(x\right)f_2\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^{r+1}dx\end{aligned}\tag{3.9}$$

and

$$\begin{aligned}X\left(t\right)&\geq\left(\tfrac{k_1}{2}-\tfrac{1}{2\epsilon_0\epsilon}\right)\int_\Omega a\left(x\right)f_1\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^2dx\\&+\left(k-\tfrac{\epsilon}{\epsilon_0}\right)\int_\Omega\ln^{\beta+1}\left(b+q\left(x\right)+t\right)\left(\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx\\&+\int_\Omega a\left(x\right)f_2\left(q\left(x\right)+t\right)\left|u\left(t\right)\right|^{r+1}dx\end{aligned}\tag{3.10}$$

for all $\epsilon>0$. We choose

δ such that $C\delta\leq\frac{\delta_0}{4}$,
ϵ such that $k-\frac{\epsilon}{\epsilon_0}\geq\frac{1-\delta_0}{8(\beta+1)}$,
k_1 such that $\frac{k_1}{2}-\frac{1}{2\epsilon_0\epsilon}-\frac{k^2(\beta+1)^2}{\epsilon}\geq\delta_0$ and $\frac{k_1}{4}-\frac{2}{\epsilon_0}-C_{T,\delta}\geq\delta_0$.

Therefore using (3.8) we obtain

$$\begin{aligned}&X\left(t+T\right)-X\left(t\right)+\tfrac{\delta_0}{4}\int_t^{t+T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&-\delta_0\int_t^{t+T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds-\tfrac{1}{2}\int_t^{t+T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds\\&+\tfrac{1-\delta_0}{8(\beta+1)}\int_t^{t+T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\\&\leq k_3\int_t^{t+T}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\end{aligned}\tag{3.11}$$

for every $t \geq T$ and for some $k_3 > 0$. Thus

$$\begin{array}{l} X\left(nT\right) + \frac{\delta_0}{4}\int_T^{nT}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2 + \left|\partial_t u\right|^2\right)dxds \\ -\delta_0\int_T^{nT}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2 dxds - \frac{1}{2}\int_T^{nT}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds \\ +\frac{1-\delta_0}{8(\beta+1)}\int_T^{nT}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds \\ \leq k_3\int_T^{nT}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left(\left|\partial_t u\right|^2 + \left|\partial_t u\right|^{2r}\right)dxds + X\left(T\right),\ \text{for all } n \geq 1. \end{array} \tag{3.12}$$

Using proposition 1, we deduce that

$$\begin{array}{l} X\left(T\right) + \frac{\delta_0}{4}\int_0^{T}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2 + \left|\partial_t u\right|^2\right)dxds \\ -\delta_0\int_0^{T}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2 dxds - \frac{1}{2}\int_0^{T}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds \\ +\frac{1-\delta_0}{8(\beta+1)}\int_0^{T}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds \leq C_T I_0 \end{array} \tag{3.13}$$

with

$$\begin{array}{l} I_0 = \left\|u_0\right\|_{H^2}^2 + \left\|u_1\right\|_{H^1}^2 + \left\|u_1\right\|_{H^1}^{2r} + \left\|u_0\right\|_{L^{r+1}}^{r+1} + \left\|\left(\ln\left(b+q\right)\right)^{\frac{\gamma}{2}}\nabla u_0\right\|_{L^2}^2 \\ +\left\|\left(\ln\left(b+q\right)\right)^{\frac{\gamma}{2}}u_1\right\|_{L^2}^2 + \left(\left\|u_0\right\|_{H^2}^2 + \left\|u_1\right\|_{H^1}^2 + \left\|u_1\right\|_{H^1}^{2r}\right)^{\frac{p}{2}} + 1 \end{array}$$

Combining (3.12) and (3.13), we obtain

$$\begin{array}{l} X\left(nT\right) + \frac{\delta_0}{4}\int_0^{nT}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2 + \left|\partial_t u\right|^2\right)dxds \\ -\delta_0\int_0^{nT}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2 dxds - \frac{1}{2}\int_0^{nT}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds \\ +\frac{1-\delta_0}{8(\beta+1)}\int_0^{nT}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds \\ \leq k_4\left(\int_0^{nT}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left(\left|\partial_t u\right|^2 + \left|\partial_t u\right|^{2r}\right)dxds + I_0\right),\ \text{for all } n \in \mathbb{N}. \end{array} \tag{3.14}$$

for some $k_4 > 0$. We remind that

$$p = \left\{\begin{array}{ll} 2\left(r+1\right) & \text{if } d \leq 3 \\ \frac{2d}{d-2} & \text{if } d \geq 4 \end{array}\right.$$

We have $r+1 < 2r < p$, using interpolation inequality and Young's inequality, we obtain

$$\begin{array}{l} \int_0^{nT}\int_\Omega a\left(x\right)f\left(q\left(x\right)+s\right)\left|\partial_t u\right|^{2r}dxds \\ \leq \int_0^{nT} f\left(s\right)\int_\Omega a\left(x\right)\left|\partial_t u\right|^{2r}dxds \\ \leq \int_0^{nT} f\left(s\right)\left(\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dx\right)^{\frac{p-2r}{p-r-1}}\left(\int_\Omega a\left(x\right)\left|\partial_t u\right|^{p}dx\right)^{\frac{r-1}{p-r-1}}ds \\ \leq \left(\left\|a\right\|_{L^\infty}\left\|\partial_t u\right\|_{L^\infty(\mathbb{R}_+,L^p(\Omega))}^p\int_0^{nT}\left(f\left(s\right)\right)^{\frac{p-r-1}{r-1}}\left(\ln\left(b+s\right)\right)^{-\frac{(\beta+1)(p-2r)}{r-1}}ds\right)^{\frac{r-1}{p-r-1}} \\ \times\left(\int_0^{nT}\ln^{\beta+1}\left(b+s\right)\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dxds\right)^{\frac{p-2r}{p-r-1}} \\ \leq \frac{\epsilon^{-\frac{p-2r}{r-1}}(r-1)\left\|a\right\|_{L^\infty}\left\|\partial_t u\right\|_{L^\infty(\mathbb{R}_+,L^p(\Omega))}^p}{p-r-1}\int_0^{+\infty}\left(b+s\right)^{-\frac{p-r-1}{r-1}}\left(\ln\left(b+s\right)\right)^{\beta-\frac{p-2r}{r-1}}ds \\ +\frac{\epsilon(p-2r)}{p-r-1}\int_0^{nT}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds \end{array}$$

for all $\epsilon > 0$. Thus using the fact that and (1.3) and Sobolev embedding $H^1 \hookrightarrow L^q$ $2 \le q \le 2 + \frac{2}{d}$, we get

$$\begin{aligned}
&\int_0^{+\infty} \int_\Omega a(x) f(q(x)+s) |\partial_t u|^{2r} dxds \\
&\le \epsilon^{-\frac{p-2r}{r-1}} C \|a\|_{L^\infty} \left(\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r} \right)^{\frac{p}{2}} \\
&+ \tfrac{\epsilon(p-2r)}{p-r-1} \int_0^{nT} \int_\Omega a(x) (1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1} dxds.
\end{aligned}$$

for all $\epsilon > 0$. To estimate the last term we have to use Holder's inequality

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x) f(q(x)+s) |\partial_t u|^2 dxds \\
&\le \|a\|_{L^\infty}^{\frac{r-1}{r+1}} \left(\int_0^{nT} \int_\Omega (f(q(x)+s))^{\frac{r+1}{r-1}} \ln^{-\frac{2(\beta+1)}{r-1}} (b+q(x)+s) dxds \right)^{\frac{r-1}{r+1}} \\
&\times \left(\int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds \right)^{\frac{2}{r+1}} \\
&\le \left(\|a\|_{L^\infty} \int_0^{+\infty} \int_\Omega (b+q(x)+s)^{-\frac{r+1}{r-1}} \ln^{\beta-\frac{2}{r-1}} (b+q(x)+s) dxds \right)^{\frac{r-1}{r+1}} \\
&\times \left(\int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds \right)^{\frac{2}{r+1}}
\end{aligned}$$

By Young's inequality, we get

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x) f(q(x)+s) |\partial_t u|^2 dxds \\
&\le \tfrac{(r-1)\epsilon^{-\frac{2}{r-1}} \|a\|_{L^\infty}}{r+1} \int_0^{+\infty} \int_\Omega (b+q(x)+s)^{-\frac{r+1}{r-1}} \ln^{\beta-\frac{2}{r-1}} (b+q(x)+s) dxds \\
&+ \tfrac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds \\
&\le C \|a\|_{L^\infty} \tfrac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \int_0^{+\infty} \ln^{\beta-\frac{2}{r-1}} (b+s) \int_0^{+\infty} (b+y+s)^{-\frac{r+1}{r-1}+d-1} dyds \\
&+ \tfrac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds \\
&\le C \|a\|_{L^\infty} \tfrac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \int_0^{+\infty} \ln^{\beta-\frac{2}{r-1}} (b+s) (b+s)^{-\frac{r+1}{r-1}+d-1} ds \\
&+ \tfrac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds
\end{aligned}$$

for all $\epsilon > 0$. Using the fact that

$$\begin{array}{ll}
-\frac{r+1}{r-1} + d < -1 & \text{if } 1 < r < 1 + \frac{2}{d} \\
\beta - \frac{2}{r-1} < -1 \text{ and } -\frac{r+1}{r-1} + d = -1, & \text{if } r = 1 + \frac{2}{d}
\end{array} \tag{3.15}$$

we get

$$\begin{aligned}
&\int_0^{+\infty} \int_\Omega a(x) f(q(x)+s) |\partial_t u|^2 dxds \\
&\le C\epsilon^{-\frac{2}{r-1}} \|a\|_{L^\infty} + \tfrac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x) \ln^{\beta+1} (b+q(x)+s) |\partial_t u|^{r+1} dxds
\end{aligned}$$

for all $\epsilon > 0$. We choose ϵ such that

$$\frac{1-\delta_0}{8(\beta+1)} - k_4 \epsilon \left(\frac{p-2r}{p-r-1} + \frac{2}{r+1} \right) \ge \frac{1-\delta_0}{16(\beta+1)}$$

We conclude that there exists a positive constant C_1 such that

$$\begin{aligned}&X\left(nT\right)+\tfrac{\delta_0}{4}\int_0^{nT}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&-\delta_0\int_0^{nT}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds-\tfrac{1}{2}\int_0^{nT}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds\\&+\tfrac{1-\delta_0}{16(\beta+1)}\int_0^{nT}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\leq C_1I_0,\ \text{for all } n\in\mathbb{N}.\end{aligned}$$

So we obtain

$$\begin{aligned}&\sup_{\mathbb{R}_+}X\left(t\right)+\tfrac{\delta_0}{4}\int_0^{+\infty}\int_\Omega f\left(q\left(x\right)+s\right)\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&-\delta_0\int_0^{+\infty}\int_\Omega a\left(x\right)f_1'\left(q\left(x\right)+s\right)\left|u\right|^2dxds-\tfrac{1}{2}\int_0^{+\infty}\int_\Omega a\left(x\right)f_2'\left(q\left(x\right)+s\right)\left|u\right|^{r+1}dxds\\&+\tfrac{1-\delta_0}{16(\beta+1)}\int_0^{+\infty}\int_\Omega a\left(x\right)\ln^{\beta+1}\left(b+q\left(x\right)+s\right)\left|\partial_t u\right|^{r+1}dxds\leq C_0I_0.\end{aligned}\tag{3.16}$$

for some positive constant C_0.

4. Proof of Theorem 2

4.1. **Preliminary results.**

Proposition 4. *We assume that Hyp A holds and* (ω,T) *geometrically controls* Ω*. Let* $\delta, R_0>0$ *and* $-1\leq\beta\leq0$*. There exists* $C_{T,\delta}=C\left(T,\delta,R_0,\beta\right)>0$ *and* $T_1>0$*, such that the following inequality*

$$\begin{aligned}&\int_t^{t+T}\int_{\Omega\cap B_{R_0}}\left(1+q\left(x\right)+s\right)^\beta\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&\leq C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^\beta\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\\&+C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-2}\left|u\right|^2dxds\\&+\delta\int_t^{t+T}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds,\end{aligned}\tag{4.1}$$

holds for every $t\geq T_1$ *and for all* u *solution of* (1.1) *with initial data* (u_0,u_1) *in* $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right).$

Proof. We set

$$\varphi\left(s\right)=\left(1+s\right)^\beta$$

It is clear that there exists a positive constant K, such that

$$\sup_{\mathbb{R}_+}\left|\frac{\varphi''(t)}{\varphi'(t)}\right|\leq K.$$

In addition the function $t\longmapsto\left|\frac{\varphi'(t)}{\varphi(t)}\right|$ is decreasing and $\lim\limits_{t\to+\infty}\left|\frac{\varphi'(t)}{\varphi(t)}\right|=0$. Moreover there exists $C>0$, such that

$$\frac{(\varphi'(t))^2}{\varphi(t)}\leq C\ \left(-\varphi'\left(t\right)\right),\ \text{for all } t\geq0.$$

Since

$$\partial_t u\in L^\infty\left(\mathbb{R}_+,H_0^1\left(\Omega\right)\right).$$

From Sobolev imbedding, we deduce that

$$\sqrt{a\varphi}\left|\partial_t u\right|^r\in L_{loc}^2\left(\mathbb{R}_+,L^2\left(\Omega\right)\right).$$

Let $\chi \in C_0^\infty\left(\mathbb{R}^d\right)$ such that

$$\chi(x) = \left\{ \begin{array}{ll} 1 & \text{for } |x| \leq R_0 \\ 0 & \text{for } |x| \geq 2R_0 \end{array} \right. .$$

Therefore using (2.17) , we infer that

$$\begin{array}{l} \int_t^{t+T} \int_{\Omega \cap B_{R_0}} (1+q(x)+s)^\beta \left(|u|^2 + |\nabla u|^2 + |\partial_t u|^2\right) dxds \\ \leq C_{T,\delta} \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^\beta \left(|\partial_t u|^2 + |\partial_t u|^{2r}\right) dxds \\ +C_{T,\delta} \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^{\beta-2} |u|^2 dxds \\ +\delta \int_t^{t+T} \int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds, \end{array} \tag{4.2}$$

This finishes the proof of the proposition. □

In order to prove theorem 2 we need the following result on the auxiliary functional X.

Lemma 2. *Let $0 < \delta_0 \ll \frac{1}{2}$. Setting $w = \psi u$ and $v = (1-\psi)u$ where u is a solution of (1.1) with initial data in $H_0^1(\Omega) \cap H^2(\Omega) \times H_0^1(\Omega)$ such that*

$$\left\|(1+q)^{\frac{\gamma}{2}} \nabla u_0\right\|_{L^2}^2 + \left\|(1+q)^{\frac{\gamma}{2}} u_1\right\|_{L^2}^2 < +\infty.$$

Let

$$\begin{array}{l} X(t) = \int_\Omega (1+q(x)+t)^\beta v(t)\,\partial_t v(t)\,dx + \frac{k_1}{2} \int_\Omega (1+q(x)+t)^{\beta-1} a(x)|u(t)|^2 dx \\ +k_2 \int_\Omega a(x)(1+q(x)+t)^{\beta-r+1} |u(t)|^{r+1} dx + \frac{k}{2} \int_\Omega (1+q(x)+t)^{\beta+1} \left(|\nabla u|^2 + |\partial_t u|^2\right) dx \end{array}$$

where

$$\begin{array}{l} k = \frac{5(1+\delta_0)r^2 + 8\left(\frac{1}{2}-\delta_0\right)(r+1) + 8\left(\frac{8}{3}\right)^r (1+\delta_0)}{10r(r+1)} \\ + \frac{\sqrt{\left(5r^2(1+\delta_0) + 8(r+1)\left(\frac{1}{2}-\delta_0\right) + 8\left(\frac{8}{3}\right)^r (1+\delta_0)\right)^2 - 160\left((1+\delta_0)\left(\frac{1}{2}-\delta_0\right)\right) r^2 (r+1)}}{10r(r+1)} \end{array}$$

and

$$\begin{array}{l} k_2 = \frac{8k(1+\delta_0)}{(r+1)(5kr - 8(1-\delta_0))} \\ 0 < 1+\beta \leq \frac{1-2\delta_0}{2k} \\ k_1 > 0 \end{array}$$

We have

$$\begin{array}{l} X(t+T) - X(t) + \delta_0 \int_t^{t+T} \int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds \\ + \left(\frac{k_1}{4} - \frac{\beta(\beta-1)}{2\epsilon_0}\right) \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^{\beta-2} |u|^2 dxds \\ + \frac{\delta_0}{r+1} \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^{\beta-r} |u|^{r+1} dxds \\ + \frac{\delta_0 r}{r+1} \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1} dxds \\ \leq C \int_t^{t+T} \int_{\Omega \cap B_{2L}} (1+q(x)+s)^\beta \left(|u|^2 + |\nabla u|^2 + |\partial_t u|^2\right) dxds \\ +2\left(\frac{1}{\epsilon_0} + \frac{4}{\epsilon_0^2 k_1} + 4k_1\right) \int_t^{t+T} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^2 dxds. \end{array} \tag{4.3}$$

for some $C > 0$.

Proof. We have

$$\int_\Omega \varphi\left(q\left(x\right)\right)\left(\left|\nabla u_0\right|^2+\left|u_1\right|^2\right)dx<\infty.$$

Then from (2.2), we infer

$$\begin{aligned}&E_\varphi\left(u\right)\left(t+T\right)+\textstyle\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta+1}\left|\partial_t u\right|^{r+1}dxds\\ &\le E_\varphi\left(u\right)\left(t\right)+\left(\beta+1\right)\textstyle\int_t^{t+T}\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta}\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds.\end{aligned}\tag{4.4}$$

Setting

$$\begin{aligned}&X_0\left(t\right)=\textstyle\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta}v\left(t\right)\partial_t v\left(t\right)dx+\frac{k_1}{2}\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta-1}a\left(x\right)\left|u\left(t\right)\right|^2dx\\ &+k_2\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-r+1}\left|u\left(t\right)\right|^{r+1}dx\end{aligned}$$

We have $v=\left(1-\psi\right)u$. Then v is a solution of

$$\left\{\begin{array}{ll}\partial_t^2 v-\Delta v+a\left(x\right)\left|\partial_t u\right|^{r-1}\partial_t v=f\left(t,x\right) & \mathbb{R}_+\times\Omega,\\ v=0 & \mathbb{R}_+\times\Gamma,\\ \left(v\left(0\right),\partial_t v\left(0\right)\right)=\left(1-\psi\right)\left(u_0,u_1\right), & \end{array}\right.\tag{4.5}$$

with

$$f\left(t,x\right)=2\nabla\psi\nabla u+u\Delta\psi.$$

Using the fact that v is a solution of (4.5), we deduce that

$$\begin{aligned}&\tfrac{d}{dt}X_0\left(t\right)=\textstyle\int_\Omega\left(\left|\partial_t v\left(t\right)\right|^2-\left|\nabla v\left(t\right)\right|^2-a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r-1}v\partial_t v\left(t\right)+f\left(t,x\right)v\right)\left(1+q\left(x\right)+t\right)^{\beta}dx\\ &+\beta\textstyle\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta-1}v\left(t\right)\partial_t v\left(t\right)dx+\frac{\beta\left(\beta-1\right)}{2}\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta-2}\left|v\right|^2dx\\ &+\tfrac{\beta}{2}\textstyle\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta-1}\left(\frac{d}{q}-\frac{\left|x\right|^2}{q^3}\right)\left|v\right|^2dx\\ &k_1\left(\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-1}u\left(t\right)\partial_t u\left(t\right)dx+\frac{\beta-1}{2}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-2}\left|u\right|^2dx\right)\\ &k_2\left(\left(\beta-r+1\right)\textstyle\int_\Omega a\left(1+q\left(x\right)+t\right)^{\beta-r}\left|u\right|^{r+1}dx+\left(r+1\right)\int_\Omega a\left(1+q\left(x\right)+t\right)^{\beta-r+1}\left|u\right|^{r-1}u\partial_t udx\right)\end{aligned}\tag{4.6}$$

By taking into account that the support of $\left(1-\psi\right)$ is contained in the set $\left\{x\in\Omega,a\left(x\right)>\epsilon_0\right\}$ and using Young's inequality we infer that

$$\begin{aligned}&\left|\beta\textstyle\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta-1}v\left(t\right)\partial_t v\left(t\right)dx\right|\\ &\le\tfrac{k_1}{8}\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-2}\left|u\left(t\right)\right|^2dx+\frac{8\beta^2}{\epsilon_0^2k_1}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta}\left|\partial_t u\left(t\right)\right|^2dx\end{aligned}$$

and

$$\begin{aligned}&k_1\left|\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-1}u\left(t\right)\partial_t u\left(t\right)dx\right|\\ &\le\tfrac{k_1}{8}\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-2}\left|u\left(t\right)\right|^2dx+8k_1\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta}\left|\partial_t u\left(t\right)\right|^2dx\end{aligned}$$

Using Young's inequality we deduce that

$$\begin{aligned}&\left|\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta}\left|\partial_t u\left(t\right)\right|^{r-1}v\partial_t v\left(t\right)dx\right|\\ &\le\tfrac{r}{r+1}\textstyle\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta+1}\left|\partial_t u\left(t\right)\right|^{r+1}dx+\frac{1}{r+1}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-r}\left|u\left(t\right)\right|^{r+1}dx\end{aligned}$$

and

$$k_2(r+1)\left|\int_\Omega a(x)(1+q(x)+t)^{\beta-r+1}|u(t)|^{r-1}u\partial_t u(t)\,dx\right|$$
$$\leq k_2\left(\tfrac{8}{3}\right)^r\int_\Omega a(x)(1+q(x)+t)^{\beta+1}|\partial_t u(t)|^{r+1}dx+\tfrac{3k_2 r}{8}\int_\Omega a(x)(1+q(x)+t)^{\beta-r}|u(t)|^{r+1}dx$$

Since $a(x)>\epsilon_0$ on the support of $(1-\psi)$, therefore

$$\begin{aligned}&\int_\Omega(1+q(x)+t)^\beta\left(|\partial_t v|^2-|\nabla v|^2\right)dx\\&=2\int_\Omega(1+q(x)+t)^\beta|\partial_t v|^2dx-\int_\Omega(1+q(x)+t)^\beta\left(|\nabla v|^2+|\partial_t v|^2\right)dx\\&\leq 2\int_\Omega(1+q(x)+t)^\beta|\partial_t v|^2dx-\tfrac{1}{2}\int_\Omega(1+q(x)+t)^\beta\left(|\nabla u|^2+|\partial_t u|^2\right)dx\\&+\int_\Omega(1+q(x)+t)^\beta\left(|\nabla w|^2+|\partial_t w|^2\right)dx\\&\leq\tfrac{2}{\epsilon_0}\int_\Omega a(x)(1+q(x)+t)^\beta|\partial_t u|^2dx-\tfrac{1}{2}\int_\Omega(1+q(x)+t)^\beta\left(|\nabla u|^2+|\partial_t u|^2\right)dx\\&+\int_\Omega(1+q(x)+t)^\beta\left(|\nabla w|^2+|\partial_t w|^2\right)dx\end{aligned}$$

Since $\psi\in C_0^\infty\left(\mathbb{R}^d\right)$ and support of ψ is contained in B_R. It easy to see that

$$\left|\int_\Omega(1+q(x)+t)^\beta f(t,x)\,v\,dx\right|\leq C\int_{\Omega\cap B_{2L}}(1+q(x)+t)^\beta\left(|u|^2+|\nabla u|^2\right)dx$$

We have $w=\psi u$. Therefore there exists a positive constant C such that

$$\begin{aligned}&\int_t^{t+T}\int_\Omega(1+q(x)+s)^\beta\left(|\nabla w(s)|^2+|\partial_t w(s)|^2\right)dxds\\&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}}(1+q(x)+s)^\beta\left(|u|^2+|\nabla u|^2+|\partial_t u|^2\right)dxds\end{aligned}$$

By taking into account of the estimates above and integrating (4.6) between t and $t+T$, we obtain

$$\begin{aligned}&X(t+T)-X(t)+\left(\tfrac{1}{2}-(1+\beta)k\right)\int_t^{t+T}\int_\Omega(1+q(x)+s)^\beta\left(|\nabla u|^2+|\partial_t u|^2\right)dxds\\&-\tfrac{\beta(d-1)}{2}\int_t^{t+T}\int_\Omega(1+q(x)+s)^{\beta-1}\tfrac{|v(t)|^2}{q}dxds\\&+\left(\tfrac{k_1}{4}-\tfrac{\beta(\beta-1)}{2\epsilon_0}\right)\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u|^2dxds\\&+\left(k_2\left(\tfrac{5}{8}r-\beta-1\right)-\tfrac{1}{r+1}\right)\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta-r}|u|^{r+1}dxds\\&+\left(k-\tfrac{r}{r+1}-k_2\left(\tfrac{8}{3}\right)^r\right)\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1}dxds\\&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}}(1+q(x)+s)^\beta\left(|u|^2+|\nabla u|^2+|\partial_t u|^2\right)dxds\\&+\left(\tfrac{2}{\epsilon_0}+\tfrac{8\beta^2}{\epsilon_0^2k_1}+8k_1\right)\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^\beta|\partial_t u|^2dxds.\end{aligned}$$

We remind that

$$\frac{1}{2}-(1+\beta)k\geq\delta_0$$

Using the fact that $k_2=\frac{8k(1+\delta_0)}{(r+1)\left(5kr-8\left(\frac{1}{2}-\delta_0\right)\right)}$, we infer that

$$k_2\left(\tfrac{5}{8}r-\beta-1\right)-\tfrac{1}{r+1}\geq k_2\left(\tfrac{5}{8}r-\tfrac{\frac{1}{2}-\delta_0}{k}\right)-\tfrac{1}{r+1}=\tfrac{\delta_0}{r+1}$$

Now we estimate the term $k-\frac{r}{r+1}-k_2\left(\frac{8}{3}\right)^r$.

$$\begin{aligned}
&k-\tfrac{(1+\delta_0)r}{r+1}-k_2\left(\tfrac{8}{3}\right)^r\\
&=k-\tfrac{(1+\delta_0)r}{r+1}-\tfrac{8k(1+\delta_0)\left(\frac{8}{3}\right)^r}{(r+1)\left(5kr-8\left(\frac{1}{2}-\delta_0\right)\right)}\\
&=\tfrac{5k^2r(r+1)-k\left(5r^2(1+\delta_0)+8(r+1)\left(\frac{1}{2}-\delta_0\right)+8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)+8\left((1+\delta_0)\left(\frac{1}{2}-\delta_0\right)\right)r}{(r+1)\left(5kr-8\left(\frac{1}{2}-\delta_0\right)\right)}=0
\end{aligned}$$

thus

$$k-\frac{r}{r+1}-k_2\left(\frac{8}{3}\right)^r=\frac{\delta_0 r}{r+1}.$$

Which yields

$$\begin{aligned}
&X(t+T)-X(t)+\delta_0\textstyle\int_t^{t+T}\int_\Omega(1+q(x)+s)^\beta\left(|\nabla u|^2+|\partial_t u|^2\right)dxds\\
&+\left(\tfrac{k_1}{4}-\tfrac{\beta(\beta-1)}{2\epsilon_0}\right)\textstyle\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u(t)|^2dxds\\
&+\tfrac{\delta_0}{r+1}\textstyle\int_t^{t+T}\int_\Omega a(1+q(x)+s)^{\beta-r}|u|^{r+1}dxds+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1}dxds\\
&\le C\textstyle\int_t^{t+T}\int_{\Omega\cap B_{2L}}(1+q(x)+s)^\beta\left(|u|^2+|\nabla u|^2+|\partial_t u|^2\right)dxds\\
&+\left(\tfrac{2}{\epsilon_0}+\tfrac{8}{\epsilon_0^2k_1}+8k_1\right)\textstyle\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^\beta|\partial_t u|^2dxds.
\end{aligned}$$

□

4.2. **Proof of Theorem 2.** We assume that Hyp A holds and (ω,T_0) geometrically controls Ω. Let u be a solution of (1.1) with initial data in $H_0^1(\Omega)\cap H^2(\Omega)\times H_0^1(\Omega)$ such that

$$\left\|(1+q)^{\frac{\gamma}{2}}\nabla u_0\right\|_{L^2}^2+\left\|(1+q)^{\frac{\gamma}{2}}u_1\right\|_{L^2}^2<+\infty.$$

Let $0<\delta_0\ll\frac{1}{2}$ and $T\ge\max(T_0,T_1)$. We define

$$\begin{aligned}
&X(t)=\textstyle\int_\Omega(1+q(x)+t)^\beta v(t)\partial_t v(t)dx+\frac{k_1}{2}\int_\Omega(1+q(x)+t)^{\beta-1}a(x)|u(t)|^2dx\\
&+k_2\textstyle\int_\Omega a(x)(1+q(x)+t)^{\beta-r+1}|u(t)|^{r+1}dx+\frac{k}{2}\int_\Omega(1+q(x)+t)^{\beta+1}\left(|\nabla u|^2+|\partial_t u|^2\right)dx,
\end{aligned}$$

where $v=(1-\psi)u$,

$$\begin{aligned}
k=&\tfrac{5(1+\delta_0)r^2+8\left(\frac{1}{2}-\delta_0\right)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)}{10r(r+1)}\\
&+\tfrac{\sqrt{\left(5r^2(1+\delta_0)+8(r+1)\left(\frac{1}{2}-\delta_0\right)+8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)^2-160\left((1+\delta_0)\left(\frac{1}{2}-\delta_0\right)\right)r^2(r+1)}}{10r(r+1)}
\end{aligned}$$

and

$$\begin{aligned}
&k_2=\tfrac{8k(1+\delta_0)}{(r+1)(5kr-8(1-\delta_0))}\\
&0<\gamma=1+\beta<\min\left(\tfrac{\frac{1}{2}-\delta_0}{k},\tfrac{d+2-dr}{r-1},\tfrac{p-2r}{r-1}\right)\\
&k_1>0.
\end{aligned}$$

According to lemma 2,

$$\begin{aligned}
&X\left(t+T\right)-X\left(t\right)+\delta_0\int_t^{t+T}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&+\left(\tfrac{k_1}{4}-\tfrac{\beta(\beta-1)}{2\epsilon_0}\right)\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-2}\left|u\right|^2dxds\\
&+\tfrac{\delta_0}{r+1}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-r}\left|u\right|^{r+1}dxds\\
&+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta+1}\left|\partial_t u\right|^{r+1}dxds\\
&\leq C\int_t^{t+T}\int_{\Omega\cap B_{2L}}\left(1+q\left(x\right)+s\right)^\beta\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dx\\
&+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^\beta\left|\partial_t u\right|^2dxds.
\end{aligned}\tag{4.7}$$

The estimate (4.1), gives

$$\begin{aligned}
&C\int_t^{t+T}\int_{\Omega\cap B_{2L}}\left(1+q\left(x\right)+s\right)^\beta\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&\leq C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^\beta\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\\
&+C_{T,\delta}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-2}\left|u\right|^2dxds\\
&+C\delta\int_t^{t+T}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds,
\end{aligned}$$

for all $t\geq T$. (4.3) and the estimates above gives

$$\begin{aligned}
&X\left(t+T\right)-X\left(t\right)+\left(\delta_0-C\delta\right)\int_t^{t+T}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\
&+\left(\tfrac{k_1}{4}-\tfrac{\beta(\beta-1)}{2\epsilon_0}-C_{T,\delta}\right)\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-2}\left|u\right|^2dxds\\
&+\tfrac{\delta_0}{r+1}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-r}\left|u\right|^{r+1}dxds\\
&+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta+1}\left|\partial_t u\right|^{r+1}dxds\\
&\leq 2\left(\tfrac{1}{\epsilon_0}+\tfrac{4}{\epsilon_0^2k_1}+4k_1+C_{T,\delta}\right)\int_t^{t+T}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^\beta\left(\left|\partial_t u\right|^{2r}+\left|\partial_t u\right|^2\right)dxds
\end{aligned}\tag{4.8}$$

for all $t\geq T$. Using Young's inequality we get

$$\begin{aligned}
&X\left(t\right)\leq\left(\tfrac{k_1}{2}+\tfrac{1}{2\epsilon_0\epsilon}\right)\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-1}\left|u\left(t\right)\right|^2dx\\
&+\left(k+\tfrac{\epsilon}{\epsilon_0}\right)\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta+1}\left(\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx\\
&+k_2\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-r+1}\left|u\left(t\right)\right|^{r+1}dx
\end{aligned}\tag{4.9}$$

and

$$\begin{aligned}
&X\left(t\right)\geq\left(\tfrac{k_1}{2}-\tfrac{1}{2\epsilon_0\epsilon}\right)\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-1}\left|u\left(t\right)\right|^2dx\\
&+\left(k-\tfrac{\epsilon}{\epsilon_0}\right)\int_\Omega\left(1+q\left(x\right)+t\right)^{\beta+1}\left(\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx\\
&+k_2\int_\Omega a\left(x\right)\left(1+q\left(x\right)+t\right)^{\beta-r+1}\left|u\left(t\right)\right|^{r+1}dx
\end{aligned}\tag{4.10}$$

for all $\epsilon > 0$. We choose

$$\begin{aligned}&\epsilon \text{ such that } k - \tfrac{\epsilon}{\epsilon_0} \geq \delta_0\\ &\delta \text{ such that } \delta_0 - C\delta \geq \tfrac{\delta_0}{2}\\ &k_1 \text{ such that } \tfrac{k_1}{2} - \tfrac{1}{2\epsilon_0\epsilon} \geq \delta_0 \text{ and } \tfrac{k_1}{4} - \tfrac{\beta(\beta-1)}{2\epsilon_0} - C_{T,\delta} \geq \delta_0.\end{aligned}$$

and we set $2\left(\frac{1}{\epsilon_0} + \frac{4}{\epsilon_0^2 k_1} + 4k_1 + C_{T,\delta}\right) = k_3$. Therefore using (4.8) we obtain

$$\begin{aligned}&X(t+T) - X(t) + \tfrac{\delta_0}{2}\int_t^{t+T}\int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds\\ &+\delta_0 \int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u|^2 dxds\\ &+\tfrac{\delta_0}{r+1}\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta-r}|u|^{r+1} dxds\\ &+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1} dxds\\ &\leq k_3 \int_t^{t+T}\int_\Omega a(x)(1+q(x)+s)^\beta \left(|\partial_t u|^{2r} + |\partial_t u|^2\right) dxds\end{aligned}$$

for all $t \geq T_1$. Thus

$$\begin{aligned}&X(nT) + \tfrac{\delta_0}{2}\int_T^{nT}\int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds\\ &+\delta_0 \int_T^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u|^2 dxds\\ &+\tfrac{\delta_0 r}{r+1}\int_T^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1} dxds\\ &+\tfrac{\delta_0}{r+1}\int_T^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta-r}|u|^{r+1} dxds\\ &\leq k_3 \int_T^{nT}\int_\Omega a(x)(1+q(x)+s)^\beta \left(|\partial_t u|^{2r} + |\partial_t u|^2\right) dxds + X(T), \text{ for all } n \geq 1.\end{aligned} \tag{4.11}$$

Using proposition 1, we obtain,

$$\begin{aligned}&X(T) + \tfrac{\delta_0}{2}\int_0^T\int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds\\ &+\delta_0 \int_0^T\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u|^2 dxds\\ &+\tfrac{\delta_0 r}{r+1}\int_0^T\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1} dxds\\ &+\tfrac{\delta_0}{r+1}\int_0^T\int_\Omega a(x)(1+q(x)+s)^{\beta-r}|u|^{r+1} dxds \leq C_T I_1.\end{aligned}$$

with

$$\begin{aligned}I_1 = &\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r} + \|u_0\|_{L^{r+1}}^{r+1} + \left\|(b+q)^{\frac{\gamma}{2}}\nabla u_0\right\|_{L^2}^2\\ &+ \left\|(1+q)^{\frac{\gamma}{2}} u_1\right\|_{L^2}^2 + \left(\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r}\right)^{\frac{p}{2}} + 1\end{aligned}$$

Therefore there exists a positive constant k_4, such that

$$\begin{aligned}&X(nT) + \tfrac{\delta_0}{2}\int_0^{nT}\int_\Omega (1+q(x)+s)^\beta \left(|\nabla u|^2 + |\partial_t u|^2\right) dxds\\ &+\delta_0 \int_0^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta-2}|u|^2 dxds\\ &+\tfrac{\delta_0 r}{r+1}\int_0^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta+1}|\partial_t u|^{r+1} dxds\\ &+\tfrac{\delta_0}{r+1}\int_0^{nT}\int_\Omega a(x)(1+q(x)+s)^{\beta-r}|u|^{r+1} dxds\\ &\leq k_4 \left(\int_0^{nT}\int_\Omega a(x)(1+q(x)+s)^\beta \left(|\partial_t u|^2 + |\partial_t u|^{2r}\right) dxds + I_1\right), \text{ for all } n \in \mathbb{N}.\end{aligned}$$

Setting $p = 2(r+1)$. We have $r + 1 < 2r < p$, using interpolation inequality and Young's inequality, we obtain

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^{2r}\, dxds \\
&\leq \int_0^{nT} (1+s)^\beta \int_\Omega a(x) |\partial_t u|^{2r}\, dxds \\
&\leq \int_0^{nT} (1+s)^\beta \left(\int_\Omega a(x) |\partial_t u|^{r+1}\, dx\right)^{\frac{p-2r}{p-r-1}} \left(\int_\Omega a(x) |\partial_t u|^{p}\, dx\right)^{\frac{r-1}{p-r-1}} ds \\
&\leq \left(\|a\|_{L^\infty} \|\partial_t u\|^p_{L^\infty(\mathbb{R}_+,L^p(\Omega))} \int_0^{+\infty} (1+s)^{\beta-\frac{2}{r-1}}\, ds\right)^{\frac{r-1}{p-r-1}} \\
&\times \left(\int_0^{nT} (1+s)^{\beta+1} \int_\Omega a(x) |\partial_t u|^{r+1}\, dxds\right)^{\frac{p-2r}{p-r-1}} \\
&\leq \frac{\epsilon^{-\frac{p-2r}{r-1}}(r-1)\|a\|_{L^\infty}\|\partial_t u\|^p_{L^\infty(\mathbb{R}_+,L^p(\Omega))}}{r+1} \int_0^{+\infty} (1+s)^{\beta-\frac{p-2r}{r-1}}\, ds \\
&+\frac{\epsilon(p-2r)}{p-r-1} \int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds
\end{aligned}$$

Now using the fact that $\beta - \frac{p-2r}{r-1} + 1 < 0$ and (1.3), we obtain

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^{2r}\, dxds \leq C\epsilon^{-\frac{p-2r}{r-1}} \|a\|_{L^\infty} \left(\|u_0\|^2_{H^2} + \|u_1\|^2_{H^1} + \|u_1\|^{2r}_{H^1}\right)^{\frac{p}{2}} \\
&+\frac{\epsilon(p-2r)}{p-r-1} \int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds.
\end{aligned}$$

To estimate the last term we use Holder's inequality

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^{2}\, dxds \\
&\leq \left(\|a\|_{L^\infty} \int_0^{nT} \int_\Omega (1+q(x)+s)^{\beta-\frac{2}{r-1}}\, dxds\right)^{\frac{r-1}{r+1}} \left(\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds\right)^{\frac{2}{r+1}}
\end{aligned}$$

By Young's inequality, we get

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^{2}\, dxds \\
&\leq \|a\|_{L^\infty} \frac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \int_0^{+\infty} \int_\Omega (1+q(x)+s)^{\beta-\frac{2}{r-1}}\, dxds \\
&+\frac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds \\
&\leq C\|a\|_{L^\infty} \frac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \int_0^{+\infty} \int_0^{+\infty} (1+y+s)^{\beta-\frac{2}{r-1}+d-1}\, dyds \\
&+\frac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds
\end{aligned}$$

Now using the fact that $\beta - \frac{2}{r-1} + d + 1 < 0$, we get

$$\begin{aligned}
&\int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^\beta |\partial_t u|^{2}\, dxds \\
&\leq C\epsilon^{-\frac{2}{r-1}} + \frac{2\epsilon}{r+1} \int_0^{nT} \int_\Omega a(x)(1+q(x)+s)^{\beta+1} |\partial_t u|^{r+1}\, dxds
\end{aligned}$$

We choose ϵ such that

$$\frac{\delta_0 r}{r+1} - k_4\epsilon\left(\frac{p-2r}{p-r-1} + \frac{2}{r+1}\right) \geq \frac{\delta_0 r}{2(r+1)}$$

So there exists a positive constant C_0 such that

$$\begin{aligned}&X\left(nT\right)+\tfrac{\delta_0}{2}\int_0^{nT}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&+\delta_0\int_0^{nT}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-2}\left|u\right|^2dxds\\&+\tfrac{\delta_0 r}{2(r+1)}\int_0^{nT}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta+1}\left|\partial_t u\right|^{r+1}dxds\\&+\tfrac{\delta_0}{r+1}\int_0^{nT}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-r}\left|u\right|^{r+1}dxds\le C_0I_1,\ \text{for all } n\in\mathbb{N}.\end{aligned}$$

This gives

$$\begin{aligned}&\sup_{\mathbb{R}_+}X\left(t\right)+\tfrac{\delta_0}{2}\int_0^{+\infty}\int_\Omega\left(1+q\left(x\right)+s\right)^\beta\left(\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&+\delta_0\int_0^{+\infty}\int_\Omega\left(1+q\left(x\right)+s\right)^{\beta-2}a\left(x\right)\left|u\left(t\right)\right|^2dxds\\&+\tfrac{\delta_0 r}{2(r+1)}\int_0^{+\infty}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta+1}\left|\partial_t u\right|^{r+1}dxds\\&+\tfrac{\delta_0}{r+1}\int_0^{+\infty}\int_\Omega a\left(x\right)\left(1+q\left(x\right)+s\right)^{\beta-r}\left|u\right|^{r+1}dxds\le C_1I_1,\end{aligned}\tag{4.12}$$

for some $C_1>0$. This finishes the proof of theorem 2.

5. Proof of Theorem 3

5.1. **Preliminary results.** As in the proof of theorem 2 we have the following weighted observability estimate.

Corollary 1. *We assume that Hyp A holds and* (ω,T) *geometrically controls* Ω. *Let* $\delta,R_0>0$, $-1\le\beta\le0$ *and* $R\ge1$. *There exists* $C_{T,\delta,R_0}>0$ *and* $T_1>0$, *such that the following inequality*

$$\begin{aligned}&\int_t^{t+T}\int_{\Omega\cap B_{R_0}}\left(R+s\right)^\beta\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&\le C_{T,\delta,R_0}\int_t^{t+T}\left(R+s\right)^\beta\int_\Omega a\left(x\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\\&\quad+C_{T,\delta,R_0}\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\right|^2dxds+\delta\int_t^{t+T}\left(R+s\right)^\beta E_u\left(s\right)ds,\end{aligned}\tag{5.1}$$

holds for every $t\ge T_1$ *and for all* u *solution of* (1.1) *with initial data* (u_0,u_1) *in* $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$.

In order to prove theorem 3 we need the following result on the auxiliary functional X.

Lemma 3. *Let* $0<\delta_0\ll1$ *and* $R\ge1$ *such that* $O\subset B_R$. *Setting* $w=\psi u$ *and* $v=\left(1-\psi\right)u$ *where* u *is a solution of* (1.1) *with initial data in* $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$ *with support contained in* B_R. *Let*

$$\begin{aligned}&X\left(t\right)=\left(R+t\right)^\beta\int_\Omega v\left(t\right)\partial_t v\left(t\right)dx+\tfrac{k_1}{2}\left(R+t\right)^{\beta-1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^2dx\\&+k_2\left(R+t\right)^{\beta-r+1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^{r+1}dx+k\left(R+t\right)^{\beta+1}E_u\left(t\right),\end{aligned}$$

where

$$\begin{aligned}k=&\tfrac{5(1+\delta_0)r^2+8(1-\delta_0)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)}{10r(r+1)}\\&+\tfrac{\sqrt{\left(5(1+\delta_0)r^2+8(1-\delta_0)(r+1)++8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)^2-160\left(1-\delta_0^2\right)r^2(r+1)}}{10r(r+1)}\end{aligned}$$

and

$$\begin{aligned} k_2 &= \frac{8k\left(1+\delta_0\right)}{\left(r+1\right)\left(5kr-8\left(1-\delta_0\right)\right)} \\ 0 &< 1+\beta \leq \frac{1-\delta_0}{k} \\ k_1 &> 0 \end{aligned}$$

We have

$$\begin{aligned} &X\left(t+T\right)-X\left(t\right)+\delta_0\int_t^{t+T}\left(R+s\right)^{\beta}E_u\left(s\right)ds+\tfrac{k_1}{4}\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\right|^2dxds \\ &+\tfrac{\delta_0}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta-r}\int_\Omega a\left(x\right)\left|u\right|^{r+1}dxds+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta+1}\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dxds \\ &\leq C_L\int_t^{t+T}\left(R+s\right)^{\beta}\int_{\Omega\cap B_{2L}}\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds \\ &+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\left(R+s\right)^{\beta}\int_\Omega a\left(x\right)\left|\partial_t u\right|^2dxds. \end{aligned} \tag{5.2}$$

for some $C_L>0$.

Proof. We have $v=\left(1-\psi\right)u$. Then v is a solution of

$$\begin{cases} \partial_t^2 v-\Delta v+a\left(x\right)\left|\partial_t u\right|^{r-1}\partial_t v=f\left(t,x\right) & \mathbb{R}_+\times\Omega, \\ v=0 & \mathbb{R}_+\times\Gamma, \\ \left(v\left(0\right),\partial_t v\left(0\right)\right)=\left(1-\psi\right)\left(u_0,u_1\right), & \end{cases} \tag{5.3}$$

with

$$f\left(t,x\right)=2\nabla\psi\nabla u+u\Delta\psi.$$

Using the fact that v is a solution of (5.3) and that

$$\frac{d}{dt}E_u\left(t\right)=-\int_\Omega a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r+1}dx,$$

we deduce that

$$\begin{aligned} &\tfrac{d}{dt}X\left(t\right)=\left(R+t\right)^{\beta}\int_\Omega\left|\partial_t v\left(t\right)\right|^2-\left|\nabla v\left(t\right)\right|^2-a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r-1}v\partial_t v\left(t\right)+f\left(t,x\right)vdx \\ &+\beta\left(R+t\right)^{\beta-1}\int_\Omega v\left(t\right)\partial_t v\left(t\right)dx \\ &k_1\left(\left(R+t\right)^{\beta-1}\int_\Omega a\left(x\right)u\left(t\right)\partial_t u\left(t\right)dx+\tfrac{\beta-1}{2}\left(R+t\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^2dx\right) \\ &k_2\left(\left(\beta-r+1\right)\left(R+t\right)^{\beta-r}\int_\Omega a\left|u\right|^{r+1}dx+\left(r+1\right)\left(R+t\right)^{\beta-r+1}\int_\Omega a\left|u\right|^{r-1}u\partial_t udx\right) \\ &+k\left(\left(\beta+1\right)\left(R+t\right)^{\beta}E_u\left(t\right)-\left(R+t\right)^{\beta+1}\int_\Omega a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r+1}dx\right). \end{aligned}$$

It easy to see that

$$\begin{aligned} \int_\Omega f\left(t,x\right)vdx &= \int_\Omega\left(2\nabla\psi\nabla u+u\Delta\psi\right)\left(1-\psi\right)udx \\ &= \int_\Omega\nabla\psi\nabla u^2+u^2\Delta\psi dx-\int_\Omega\frac{1}{2}\nabla\psi^2\nabla u^2+u^2\psi\Delta\psi dx \\ &= \int_\Omega\left|\nabla\psi\right|^2\left|u\right|^2dx. \end{aligned}$$

By taking into account that the support of $(1-\psi)$ is contained in the set $\{x\in\Omega, a(x)>\epsilon_0\}$ and using Young's inequality we infer that

$$\begin{aligned}&\left|\beta(R+t)^{\beta-1}\int_\Omega v(t)\,\partial_t v(t)\,dx\right|\\&\leq(R+t)^{\beta-2}\tfrac{k_1}{8}\int_\Omega a(x)\left|u(t)\right|^2dx+\tfrac{8\beta^2}{\epsilon_0^2k_1}(R+t)^\beta\int_\Omega a(x)\left|\partial_t u(t)\right|^2dx\end{aligned}$$

and

$$\begin{aligned}&\left|k_1(R+t)^{\beta-1}\int_\Omega au(t)\,\partial_t u(t)\,dx\right|\\&\leq(R+t)^{\beta-2}\tfrac{k_1}{8}\int_\Omega a(x)\left|u(t)\right|^2dx+8k_1(R+t)^\beta\int_\Omega a(x)\left|\partial_t u(t)\right|^2dx\end{aligned}$$

Using Young's inequality we deduce that

$$\begin{aligned}&\left|(R+t)^\beta\int_\Omega a(x)\left|\partial_t u(t)\right|^{r-1}v\partial_t v(t)\,dx\right|\\&\leq\tfrac{r(R+t)^{\beta+1}}{r+1}\int_\Omega a(x)\left|\partial_t u(t)\right|^{r+1}dx+\tfrac{(R+t)^{\beta-r}}{r+1}\int_\Omega a(x)\left|u(t)\right|^{r+1}dx\end{aligned}$$

and

$$\begin{aligned}&k_2(r+1)\left|(R+t)^{\beta-r+1}\int_\Omega a(x)\left|u(t)\right|^{r-1}u\partial_t u(t)\,dx\right|\\&\leq k_2\left(\tfrac{8}{3}\right)^r(R+t)^{\beta+1}\int_\Omega a(x)\left|\partial_t u(t)\right|^{r+1}dx+\tfrac{3k_2r(R+t)^{\beta-r}}{8}\int_\Omega a(x)\left|u(t)\right|^{r+1}dx\end{aligned}$$

Since $a(x)>\epsilon_0$ on the support of $(1-\psi)$, therefore

$$\begin{aligned}&(R+t)^\beta\int_\Omega\left|\partial_t v(t)\right|^2-\left|\nabla v(t)\right|^2dx\\&\leq\tfrac{2(R+t)^\beta}{\epsilon_0}\int_\Omega a(x)\left|\partial_t u(t)\right|^2dx-(R+t)^\beta E_u(t)+2(R+t)^\beta E_w(t)\end{aligned}$$

By taking into account of the estimates above, we obtain

$$\begin{aligned}&\tfrac{d}{dt}X(t)+(1-(1+\beta)k)(R+t)^\beta E_u(t)+\tfrac{k_1}{4}(R+s)^{\beta-2}\int_\Omega a(x)\left|u(t)\right|^2dx\\&+\left(k_2\left(\tfrac{5}{8}r-\beta-1\right)-\tfrac{1}{r+1}\right)(R+t)^{\beta-r}\int_\Omega a\left|u(t)\right|^{r+1}dx\\&+\left(k-\tfrac{r}{r+1}-k_2\left(\tfrac{8}{3}\right)^r\right)(R+t)^{\beta+1}\int_\Omega a(x)\left|\partial_t u(t)\right|^{r+1}dx\\&\leq2(R+t)^\beta E_w(t)+\left(\tfrac{2}{\epsilon_0}+\tfrac{8\beta^2}{\epsilon_0^2k_1}+8k_1\right)(R+t)^\beta\int_\Omega a(x)\left|\partial_t u(t)\right|^2dx\\&+(R+t)^\beta\int_\Omega\left|\nabla\psi\right|^2\left|u(t)\right|^2dx.\end{aligned}$$

We remind that

$$1-(1+\beta)k\geq\delta_0$$

Using the fact that $k_2=\frac{8k(1+\delta_0)}{(r+1)(5kr-8(1-\delta_0))}$, we infer that

$$k_2\left(\tfrac{5}{8}r-\beta-1\right)-\tfrac{1}{r+1}\geq k_2\left(\tfrac{5}{8}r-\tfrac{1-\delta_0}{k}\right)-\tfrac{1}{r+1}=\tfrac{\delta_0}{r+1}$$

Now we estimate the term $k-\frac{r}{r+1}-k_2\left(\frac{8}{3}\right)^r$. Since $k\geq k_0$, therefore

$$\begin{aligned}&k-\tfrac{(1+\delta_0)r}{r+1}-k_2\left(\tfrac{8}{3}\right)^r\\&\geq k-\tfrac{(1+\delta_0)r}{r+1}-\tfrac{8k(1+\delta_0)\left(\frac{8}{3}\right)^r}{(r+1)(5kr-8(1-\delta_0))}\\&=\tfrac{5k^2r(r+1)-k\left(5r^2(1+\delta_0)+8(r+1)(1-\delta_0)+8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)+8\left(1-\delta_0^2\right)r}{(r+1)(5k_0r-8(1-\delta_0))}=0\end{aligned}$$

thus

$$k-\frac{r}{r+1}-k_2\left(\frac{8}{3}\right)^r\geq\frac{\delta_0 r}{r+1}.$$

Which yields

$$\begin{aligned}
&\tfrac{d}{dt}X\left(t\right)+\delta_0\left(R+t\right)^{\beta}E_u\left(t\right)+\tfrac{k_1}{4}\left(R+s\right)^{\beta-2}\textstyle\int_{\Omega}a\left(x\right)\left|u\left(t\right)\right|^2dx\\
&+\tfrac{\delta_0}{r+1}\left(R+t\right)^{\beta-r}\textstyle\int_{\Omega}a\left|u\left(t\right)\right|^{r+1}dx+\tfrac{\delta_0 r}{r+1}\left(R+t\right)^{\beta+1}\textstyle\int_{\Omega}a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r+1}dx\\
&\leq 2\left(R+t\right)^{\beta}E_w\left(t\right)+\left(\tfrac{2}{\epsilon_0}+\tfrac{8}{\epsilon_0^2 k_1}+8k_1\right)\left(R+t\right)^{\beta}\textstyle\int_{\Omega}a\left(x\right)\left|\partial_t u\left(t\right)\right|^2dx\\
&+\left(R+t\right)^{\beta}\textstyle\int_{\Omega}\left|\nabla\psi\right|^2\left|u\left(t\right)\right|^2dx.
\end{aligned}\tag{5.4}$$

We have $w=\psi u$. It is clear that

$$2\left(R+t\right)^{\beta}E_w\left(t\right)ds\leq C_L\left(R+t\right)^{\beta}\textstyle\int_{\Omega\cap B_{2L}}\left(\left|u\left(t\right)\right|^2+\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx$$

in addition we have

$$\textstyle\int_{\Omega}\left|\nabla\psi\right|^2\left|u\left(t\right)\right|^2dx\leq C_L\textstyle\int_{\Omega\cap B_{2L}}\left|u\left(t\right)\right|^2dx$$

Therefore

$$\begin{aligned}
&\tfrac{d}{dt}X\left(t\right)+\delta_0\left(R+t\right)^{\beta}E_u\left(t\right)+\tfrac{k_1}{4}\left(R+t\right)^{\beta-2}\textstyle\int_{\Omega}a\left(x\right)\left|u\left(t\right)\right|^2dx\\
&+\tfrac{\delta_0}{r+1}\left(R+t\right)^{\beta-r}\textstyle\int_{\Omega}a\left|u\left(t\right)\right|^{r+1}dx+\tfrac{\delta_0 r}{r+1}\left(R+t\right)^{\beta+1}\textstyle\int_{\Omega}a\left(x\right)\left|\partial_t u\left(t\right)\right|^{r+1}dx\\
&\leq C_L\left(R+t\right)^{\beta}\textstyle\int_{\Omega\cap B_{2L}}\left(\left|u\left(t\right)\right|^2+\left|\nabla u\left(t\right)\right|^2+\left|\partial_t u\left(t\right)\right|^2\right)dx\\
&+\left(\tfrac{2}{\epsilon_0}+\tfrac{8}{\epsilon_0^2 k_1}+8k_1\right)\left(R+t\right)^{\beta}\textstyle\int_{\Omega}a\left(x\right)\left|\partial_t u\left(t\right)\right|^2dx
\end{aligned}$$

Integrating the estimate above between t and $t+T$ we get (5.2). □

5.2. **Proof of Theorem 3.** We assume that Hyp A holds and (ω,T_0) geometrically controls Ω. Let $R\geq 1$ such that $O\subset B_R$ and u be the solution of (1.1) with initial data in $H_0^1\left(\Omega\right)\cap H^2\left(\Omega\right)\times H_0^1\left(\Omega\right)$ with support contained in B_R. Let $0<\delta_0\ll 1$ and $T\geq\max\left(T_0,T_1\right)$. We define

$$\begin{aligned}
&X\left(t\right)=\left(R+t\right)^{\beta}\textstyle\int_{\Omega}v\left(t\right)\partial_t v\left(t\right)dx+\tfrac{k_1}{2}\left(R+t\right)^{\beta-1}\textstyle\int_{\Omega}a\left(x\right)\left|u\left(t\right)\right|^2dx\\
&+k_2\left(R+t\right)^{\beta-r+1}\textstyle\int_{\Omega}a\left(x\right)\left|u\left(t\right)\right|^{r+1}dx+k\left(R+t\right)^{\beta+1}E_u\left(t\right),
\end{aligned}$$

where $v=\left(1-\psi\right)u$ and

$$\begin{aligned}
k=&\tfrac{5(1+\delta_0)r^2+8(1-\delta_0)(r+1)+8\left(\frac{8}{3}\right)^r(1+\delta_0)}{10r(r+1)}\\
&+\tfrac{\sqrt{\left(5(1+\delta_0)r^2+8(1-\delta_0)(r+1)++8\left(\frac{8}{3}\right)^r(1+\delta_0)\right)^2-160\left(1-\delta_0^2\right)r^2(r+1)}}{10r(r+1)}
\end{aligned}$$

and

$$\begin{aligned}
&k_2=\tfrac{8k(1+\delta_0)}{(r+1)(5kr-8(1-\delta_0))}\\
&0<\gamma=\beta+1<\min\left(\tfrac{1-\delta_0}{k},\tfrac{d+2-dr}{r-1},\tfrac{p-2r}{r-1}\right)\\
&k_1>0.
\end{aligned}$$

According to lemma 3,

$$\begin{aligned}&X\left(t+T\right)-X\left(t\right)+\delta_0\int_t^{t+T}\left(R+s\right)^{\beta}E_u\left(s\right)ds+\tfrac{k_1}{4}\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\right|^2dxds\\&+\tfrac{\delta_0}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta-r}\int_\Omega a\left(x\right)\left|u\right|^{r+1}dxds+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta+1}\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dxds\\&\leq C_L\int_t^{t+T}\left(R+s\right)^{\beta}\int_{\Omega\cap B_{2L}}\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&+2\left(\tfrac{1}{\epsilon_0}+\tfrac{4}{\epsilon_0^2k_1}+4k_1\right)\int_t^{t+T}\left(R+s\right)^{\beta}\int_\Omega a\left(x\right)\left|\partial_t u\right|^2dxds.\end{aligned}\tag{5.5}$$

The estimate (5.1), gives

$$\begin{aligned}&C_L\int_t^{t+T}\left(R+s\right)^{\beta}\int_{\Omega\cap B_{2L}}\left(\left|u\right|^2+\left|\nabla u\right|^2+\left|\partial_t u\right|^2\right)dxds\\&\leq C_{T,\delta}\int_t^{t+T}\left(R+s\right)^{\beta}\int_\Omega a\left(x\right)\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}dxds\\&\quad+C_{T,\delta}\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\right|^2dxds+C_L\delta\int_t^{t+T}\left(R+s\right)^{\beta}E_u\left(s\right)ds,\end{aligned}$$

for all $t\geq T$. (5.2) and the estimate above gives

$$\begin{aligned}&X\left(t+T\right)-X\left(t\right)+\left(\delta_0-C_L\delta\right)\int_t^{t+T}\left(R+s\right)^{\beta}E_u\left(s\right)ds\\&+\left(\tfrac{k_1}{4}-C_{T,\delta}\right)\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\right|^2dxds\\&+\tfrac{\delta_0}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta-r}\int_\Omega a\left(x\right)\left|u\right|^{r+1}dxds\\&+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta+1}\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dxds\\&\leq 2\left(\tfrac{1}{\epsilon_0}+\tfrac{4}{\epsilon_0^2k_1}+4k_1+C_{T,\delta}\right)\int_t^{t+T}\left(R+s\right)^{\beta}\int_\Omega a\left(x\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\end{aligned}\tag{5.6}$$

Using Young's inequality we get

$$\begin{aligned}&X\left(t\right)\leq\left(\tfrac{k_1}{2}+\tfrac{1}{2\epsilon_0\epsilon}\right)\left(R+t\right)^{\beta-1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^2dx\\&+\left(k+\tfrac{\epsilon}{\epsilon_0}\right)\left(R+t\right)^{\beta+1}E_u\left(t\right)+k_2\left(R+t\right)^{\beta-r+1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^{r+1}dx\end{aligned}\tag{5.7}$$

and

$$\begin{aligned}&X\left(t\right)\geq\left(\tfrac{k_1}{2}-\tfrac{1}{2\epsilon_0\epsilon}\right)\left(R+t\right)^{\beta-1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^2dx\\&+\left(k-\tfrac{\epsilon}{\epsilon_0}\right)\left(R+t\right)^{\beta+1}E_u\left(t\right)+k_2\left(R+t\right)^{\beta-r+1}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^{r+1}dx\end{aligned}\tag{5.8}$$

for all $\epsilon>0$. We choose

$$\begin{aligned}&\epsilon\text{ such that }k-\tfrac{\epsilon}{\epsilon_0}\geq\delta_0\\&k_1\text{ such that }\tfrac{k_1}{2}-\tfrac{1}{2\epsilon_0\epsilon}\geq\delta_0\text{ and }\tfrac{k_1}{4}-C_{T,\delta}\geq\delta_0\\&\delta\text{ such that }\delta_0-C_L\delta\geq\tfrac{\delta_0}{2}.\end{aligned}$$

Therefore using (5.6) we obtain

$$\begin{aligned}&X\left(t+T\right)-X\left(t\right)+\tfrac{\delta_0}{2}\int_t^{t+T}\left(R+s\right)^{\beta}E_u\left(s\right)ds+\delta_0\int_t^{t+T}\left(R+s\right)^{\beta-2}\int_\Omega a\left(x\right)\left|u\left(t\right)\right|^2dxds\\&+\tfrac{\delta_0}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta-r}\int_\Omega a\left(x\right)\left|u\right|^{r+1}dxds+\tfrac{\delta_0 r}{r+1}\int_t^{t+T}\left(R+s\right)^{\beta+1}\int_\Omega a\left(x\right)\left|\partial_t u\right|^{r+1}dxds\\&\leq k_3\int_t^{t+T}\left(R+s\right)^{\beta}\int_\Omega a\left(x\right)\left(\left|\partial_t u\right|^2+\left|\partial_t u\right|^{2r}\right)dxds\end{aligned}\tag{5.9}$$

for all $t \geq T$ where $k_3 = 2\left(\frac{1}{\epsilon_0} + \frac{4}{\epsilon_0^2 k_1} + 4k_1 + C_L C_{T,\delta}\right)$. Thus

$$\begin{aligned} &X(nT) + \tfrac{\delta_0}{2}\textstyle\int_T^{nT} (R+s)^{\beta} E_u(s)\,ds + \delta_0 \int_T^{nT} (R+s)^{\beta-2} \int_\Omega a(x)\,|u|^2\,dxds \\ &+\tfrac{\delta_0}{r+1}\textstyle\int_T^{nT} (R+s)^{\beta-r} \int_\Omega a(x)\,|u|^{r+1}\,dxds + \tfrac{\delta_0 r}{r+1}\int_T^{nT} (R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dxds \\ &\leq k_3 \textstyle\int_T^{nT} (R+s)^{\beta} \int_\Omega a(x)\left(|\partial_t u|^2 + |\partial_t u|^{2r}\right) dxds + X(T), \text{ for all } n \geq 1. \end{aligned} \tag{5.10}$$

It is easy to see that

$$\begin{aligned} &X(T) + \tfrac{\delta_0}{2}\textstyle\int_0^{T} (R+s)^{\beta} E_u(s)\,ds + \delta_0 \int_0^{T} (R+s)^{\beta-2} \int_\Omega a(x)\,|u|^2\,dxds \\ &+\tfrac{\delta_0}{r+1}\textstyle\int_0^{T} (R+s)^{\beta-r} \int_\Omega a(x)\,|u|^{r+1}\,dxds + \tfrac{\delta_0 r}{r+1}\int_0^{T} (R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dxds \\ &\leq C_T I_2 \end{aligned}$$

with

$$I_2 = \|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_0\|_{L^{r+1}}^{r+1} + \left(\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r}\right)^{\frac{p}{2}} + 1$$

So there exists a positive constant k_4 such that

$$\begin{aligned} &X(nT) + \tfrac{\delta_0}{2}\textstyle\int_0^{nT} (R+s)^{\beta} E_u(s)\,ds + \delta_0 \int_0^{nT} (R+s)^{\beta-2} \int_\Omega a(x)\,|u|^2\,dxds \\ &+\tfrac{\delta_0}{r+1}\textstyle\int_0^{nT} (R+s)^{\beta-r} \int_\Omega a(x)\,|u|^{r+1}\,dxds + \tfrac{\delta_0 r}{r+1}\int_0^{nT} (R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dxds \\ &\leq k_3 \left(\textstyle\int_0^{nT} (R+s)^{\beta} \int_\Omega a(x)\left(|\partial_t u|^{2r} + |\partial_t u|^2\right) dxds + I_2\right), \text{ for all } n \in \mathbb{N}. \end{aligned}$$

We remind that

$$p = \begin{cases} 2(r+1) & \text{if } d \leq 3 \\ \frac{2d}{d-2} & \text{if } d \geq 4 \end{cases}$$

We proceed as in the proof of theorem 2, we get

$$\begin{aligned} &\textstyle\int_0^{nT} (R+s)^{\beta} \int_\Omega a(x)\,|\partial_t u|^{2r}\,dxds \leq C\epsilon^{-\frac{p-2r}{r-1}} \|a\|_{L^\infty} \left(\|u_0\|_{H^2}^2 + \|u_1\|_{H^1}^2 + \|u_1\|_{H^1}^{2r}\right) \\ &+\tfrac{\epsilon(p-2r)}{p-r-1}\textstyle\int_0^{nT} \int_\Omega a(x)\,(R+s)^{\beta+1}\,|\partial_t u|^{r+1}\,dxds. \end{aligned}$$

To estimate the last term we have to use the finite speed propagation property and the fact that the support of the initial data is contained in B_R.

$$\begin{aligned} &\textstyle\int_0^{nT} (R+s)^{\beta} \int_\Omega a(x)\,|\partial_t u|^2\,dxds \\ &\leq \|a\|_{L^\infty}^{\frac{r-1}{r+1}} \textstyle\int_0^{nT} (R+s)^{\beta} \left(\int_{\Omega\cap B_{R+s}} dx\right)^{\frac{r-1}{r+1}} \left(\int_\Omega a(x)\,|\partial_t u|^{r+1}\,dx\right)^{\frac{2}{r+1}} ds \\ &\leq \textstyle\int_0^{nT} \left(\|a\|_{L^\infty} \int_{\Omega\cap B_{R+s}} (R+s)^{\beta-\frac{2}{r-1}}\,dx\right)^{\frac{r-1}{r+1}} \left((R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dx\right)^{\frac{2}{r+1}} \end{aligned}$$

By Young's inequality, we get

$$\begin{aligned} &\textstyle\int_0^{nT} (R+s)^{\beta} \int_\Omega a(x)\,|\partial_t u|^2\,dxds \\ &\leq \|a\|_{L^\infty} \tfrac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \textstyle\int_0^{+\infty} \int_{\Omega\cap B_{R+s}} (R+s)^{\beta-\frac{2}{r-1}}\,dxds + \tfrac{2\epsilon}{r+1}\int_0^{nT} (R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dxds \\ &\leq C\,\|a\|_{L^\infty} \tfrac{(r-1)\epsilon^{-\frac{2}{r-1}}}{r+1} \textstyle\int_0^{+\infty} (R+s)^{\beta-\frac{2}{r-1}+d}\,ds + \tfrac{2\epsilon}{r+1}\int_0^{nT} (R+s)^{\beta+1} \int_\Omega a(x)\,|\partial_t u|^{r+1}\,dxds \end{aligned}$$

Now using the fact that $\beta - \frac{2}{r-1} + d + 1 < 0$, we get

$$\int_0^{nT} (R+s)^{\beta} \int_{\Omega} a(x) |\partial_t u|^2 \, dxds \leq C\epsilon^{-\frac{2}{r-1}} + \frac{2\epsilon}{r+1} \int_0^{+\infty} (R+s)^{\beta+1} \int_{\Omega} a(x) |\partial_t u|^{r+1} \, dxds$$

We choose ϵ such that

$$\frac{\delta_0 r}{r+1} - k_4 \epsilon \left(\frac{2}{r+1} + \frac{p-2r}{p-r-1} \right) = \frac{\delta_0 r}{2(r+1)}$$

Therefore there exists a positive constant C_0 such that

$$\begin{aligned} & X(nT) + \frac{\delta_0}{2} \int_0^{nT} (R+s)^{\beta} E_u(s) \, ds + \delta_0 \int_0^{nT} (R+s)^{\beta-2} \int_{\Omega} a(x) |u|^2 \, dxds \\ & + \frac{\delta_0}{r+1} \int_0^{nT} (R+s)^{\beta-r} \int_{\Omega} a(x) |u|^{r+1} \, dxds + \frac{\delta_0 r}{2(r+1)} \int_0^{nT} (R+s)^{\beta+1} \int_{\Omega} a(x) |\partial_t u|^{r+1} \, dxds \\ & \leq C_0 R^{1+\beta} I_2, \text{ for all } n \in \mathbb{N}. \end{aligned}$$

This gives

$$\begin{aligned} & \sup_{\mathbb{R}_+} X(t) + \frac{\delta_0}{2} \int_0^{+\infty} (R+s)^{\beta} E_u(s) \, ds + \delta_0 \int_0^{+\infty} (R+s)^{\beta-2} \int_{\Omega} a(x) |u|^2 \, dxds \\ & + \frac{\delta_0}{r+1} \int_0^{+\infty} (R+s)^{\beta-r} \int_{\Omega} a(x) |u|^{r+1} \, dxds + \frac{\delta_0 r}{2(r+1)} \int_0^{+\infty} (R+s)^{\beta+1} \int_{\Omega} a(x) |\partial_t u|^{r+1} \, dxds \\ & \leq C_2 R^{1+\beta} I_0. \end{aligned}$$

for some $C_2 > 0$. This finishes the proof of theorem 3.

References

[1] L. Aloui, S. Ibrahim and K. Nakanishi, Exponential Energy Decay for Damped Klein–Gordon Equation with Nonlinearities of Arbitrary Growth, Communications in Partial Differential Equations, Volume 36, Issue 5, 797-818, (2010).

[2] Aloui. L, Khenissi, M. Stabilisation de l'équation des ondes dans un domaine extérieur, Rev. Mat. Iberoamerica, 28 (2002), 1–16.

[3] J. Bae, M. Nakao, Energy decay for the wave equation with boundary and localized dissipations in exterior domains, Math. Nachr. 278, No. 7–8, 771 – 783 (2005).

[4] A. Bchatnia, M. Daoulatli, Local energy decay for the wave equation with a nonlinear time dependent damping. (preprint).

[5] C. Bardos, G. Lebeau, J. Rauch, Sharp sufficient conditions for the observation, control, and stabilization of waves from the boundary. SIAM J. Control Optimization 30, No.5, 1024-1065 (1992).

[6] N. Burq, Mesures semi-classiques et mesures de défaut. (French) [Semiclassical measures and defect measures] Séminaire Bourbaki, Vol. 1996/97. Astérisque245 (1997), Exp.826, 4, pp. 167-195.

[7] N. Burq, P. Gérard, Condition nécessaire et suffisante pour la contrôlabilité exacte des ondes, Comptes Rendus de l'Académie des Sciences - Series I - Mathematics, Volume 325, Issue 7, October 1997, Pages 749-752.

[8] W. Dan, Y. Shibata, On a local energy decay of solutions of a dissipative wave equation, Funkcial. Ekvac. 38 (1995) 545–568.

[9] M. Daoulatli, Local energy decay for the nonlinear dissipative wave equation in an exterior domain, Port. Math., 64 (2007), pp. 39–65.

[10] M. Daoulatli, B. Dehman, and M. Khenissi, Local energy decay for the elastic system with nonlinear damping in an exterior domain, SIAM J. Control Optim. 48, pp. 5254-5275

[11] M. Daoulatli , Behaviors of the energy of solutions of the wave equation with damping and external force, Journal of Mathematical Analysis and Applications, Volume 389, Issue 1, 1 May 2012, Pages 205–225.

[12] M. Daoulatli, Energy decay rates for solutions of the wave equation with linear damping in exterior domain, arXiv:1203.6780v4.

[13] B. Dehman, G. Lebeau and E. Zuazua, Stabilization and control for the subcritical semilinear wave equation, Ann. Scient. Éc. Norm. Sup., 4e série, t. 36, 2003, p. 525 à 551.

[14] P. Gérard. Microlocal Defect Measures . Comm. Partial Diff. eq., 16:1762–1794, 1991.

[15] R. Ikehata, G. Todorova, B. Yordanov; Critical Exponent for Semilinear Wave Equations with Space-Dependent Potential, Funkcialaj Ekvacioj, 52 (2009), 411–435.
[16] R. Ikehata, Energy decay of solutions for the semilinear dissipative wave equations in an exterior domain, Funkcial. Ekvac. 44 (2001) 487–499.
[17] R. Ikehata, Fast decay of solutions for linear wave equations with dissipation localized near infinity in an exterior domain, Journal of Differential Equations, Volume 188, Issue 2, 1 March 2003, Pages 390-405.
[18] R. Ikehata, T. Matsuyama, L2-behaviour of solutions to the linear heat and wave equations in exterior domains, Sci. Math. Japon. 55 (2002) 33–42.
[19] R. Joly, L. Camille, Stabilization for the semilinear wave equation with geometric control condition, arXiv:1205.2459
[20] S. Kawashima, M. Nakao, K. Ono, On the decay property of solutions to the Cauchy problem of the semilinear wave equation with a dissipative term, J. Math. Soc. Jpn. 47 (1995) 617–653.
[21] M. Khenissi, Équation des ondes amorties dans un domaine extérieur. Bull. Soc. Math. France 131 (2003), no. 2, 211–228.
[22] G. Lebeau. Equation des ondes amorties . In Algebraic and geometric methods in mathematical physics: proceedings of the Kaciveli Summer School, Crimea, Ukraine, 1993, page 73. Springer, 1996.
[23] J.L.Lions and W.Strauss, Some non-linear evolution equations, Bull. Soc. Math. France, 93 (1965), 43-96.
[24] Mochizuki, Kiyoshi; Motai, Takahiro, On energy decay-nondecay problems for wave equations with nonlinear dissipative term in R^N, J. Math. Soc. Japan 47 (1995), no. 3, 405–421.
[25] Nakao, M. – Stabilization of local energy in an exterior domain for the wave equation with a localized dissipation, J. Diff. Eq., 148 (1998), 388–406.
[26] M. Nakao, Energy decay for the linear and semilinear wave equations in exterior domains with some localized dissipations, Math. Z. 238 (2001) 781–797.
[27] M. Nakao, Decay of solutions to the Cauchy problem for the Klein-Gordon equation with a localized nonlinear dissipation, Hokkaido Math. J. 27 (1998), p. 245–271.
[28] M. Nakao, Energy decay of the wave equation with a nonlinear dissipative term, Funkcial. Ekvac. 26 (1983), no. 3, 237–250.
[29] M. Nakao, Energy decay to the Cauchy problem for a generalised nonlinear Klein-Gordon equation with a nonlinear dissipative term. Int. J. Dyn. Syst. Differ. Equ. 3 (2011), no. 3, 349–362.
[30] M. Nakao; Jung, Il Hyo, Energy decay for the wave equation in exterior domains with some half-linear dissipation, Differential Integral Equations 16 (2003), no. 8, 927–948.
[31] The time decay to the Cauchy problem for semilinear dissipative wave equations, Adv. Math. Sci. Appl. 9(1999), 243-262.
[32] K. Ono, L^1 estimates for dissipative wave equations in exterior domains. J. Math. Anal. Appl. 333 (2007), no. 2, 1079–1092
[33] R. Racke, Non-homogeneous non-linear damped wave equations in unbounded domains, Math. Methods Appl. Sci. 13 (1990), 481–491.
[34] L. Tartar. H-measures, a new approach for studying homogenisation, oscillations and concentration effects in partial differential equations . Proceedings of the Royal Society of Edinburgh. Section A. Mathematics, 115 (3-4):193–230, 1990
[35] G. Todorova, B. Yordanov, The energy decay problem for wave equations with nonlinear dissipative terms in Rn. Indiana Univ. Math. J. 56 (2007), no. 1, 389–416.
[36] E. Zuazua, Exponential deacy for the semilinear wave equation with locallized damping in unbounded domains, J. Math. pures et appl. 70 (1992), p. 513–529.

Department of Mathematics, FSB, University of Carthage & LAMSIN, ENIT, University of Tunis Elmanar
E-mail address, M. Daoulatli: `moez.daoulatli@infcom.rnu.tn`